



AFAL-TR-88-056

AD:

AD-A203 801

Final Report  
for the period  
November 1985 to  
February 1988

# Condensation and Storage of Hydrogen Cluster Ions

November 1988

Author:  
J. T. Bahns

University of Dayton  
Research Institute  
300 College Park  
Dayton, OH 45469-0001

UDR-TR-88-28  
F04611-83-C-0046

## Approved for Public Release

Distribution is unlimited. The AFAL Technical Services Office has reviewed this report, and it is releasable to the National Technical Information Service, where it will be available to the general public, including foreign nationals.

Prepared for the:

Air Force  
Astronautics  
Laboratory

Air Force Space Technology Center  
Space Division, Air Force Systems Command  
Edwards Air Force Base,  
California 93523-5000

DTIC  
ELECTE  
S D  
E

89 1 18 097


## NOTICE


When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related government procurement operation, the government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

## FOREWORD


This final report was submitted by the University of Dayton in support of contract F04611-83-C-0046 with the Air Force Astronautics Laboratory (AFAL), Edwards Air Force Base, CA. AFAL Project Manager was Lt Monte Turner.

This report has been reviewed and is approved for distribution in accordance with the the distribution statement on the cover and on the DD Form 1473.

  
MONTE D. TURNER, 1Lt, USAF  
Project Manager

  
WAYNE E. ROE  
Research Coordinator

FOR THE COMMANDER

  
STEPHEN L. RODGERS  
Chief, ARIES Office

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION /AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.			
2b. DECLASSIFICATION /DOWNGRADING SCHEDULE						
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UDR-TR-88-28			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFAL-TR-88-056			
6a. NAME OF PERFORMING ORGANIZATION University of Dayton Research Institute		6b. OFFICE SYMBOL (If applicable) UDRI		7a. NAME OF MONITORING ORGANIZATION Air Force Astronautics Laboratory		
6c. ADDRESS (City, State, and ZIP Code) 300 College Park Dayton, OH 45469-0001			7b. ADDRESS (City, State, and ZIP Code) LSX Edwards AFB, CA 93523-5000			
8a. NAME OF FUNDING /SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F04611-83-C-0046		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS			
			PROGRAM ELEMENT NO 61102F	PROJECT NO. 2301	TASK NO M1	WORK UNIT ACCESSION NO RB
11. TITLE (Include Security Classification) Condensation and Storage of Hydrogen Cluster Ions (U)						
12. PERSONAL AUTHOR(S) Bahns, John T.						
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 85/11 TO 88/2		14. DATE OF REPORT (Year, Month, Day) 88/11		15. PAGE COUNT 105
16. SUPPLEMENTARY NOTATION						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP	SUB-GROUP				
07	02		CI approach hydrogen cluster ions, pseudo wall			
07	04		containerless nonneutral plasma, penning trap			
			condensation, annular plasma, hydrogen cluster.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)						
<p>This report reviews the cluster ion approach to containerless condensation and introduces the concept of pseudo wall. An experimental pseudo wall device, designed for the study of containerless condensation techniques, is described. Calculations indicate that the proposed device will be capable of confining cluster ions in the mass range 2-10<sup>7</sup> for time exceeding 10<sup>4</sup> seconds. A "next generation" device is projected to be capable of containing a total mass of 0.3 milligrams.</p>						
20. DISTRIBUTION /AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS				21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Monte D. Turner, 1Lt., USAF				22b. TELEPHONE (Include Area Code) (805) 275-5749		22c. OFFICE SYMBOL LSX

Block 18 (continued): ion plasma, classical Brillouin limit.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
NTIS DB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Classification	
By	
Distribution	
Availability Codes	
Dist	Avail and/or Special
A-1	



## SUMMARY

This report briefly reviews our current understanding of the Cluster Ion (CI) approach, describes an experimental apparatus that will (for the first time) allow long-term studies ( $>10^4$  seconds) of very large hydrogen cluster ions ( $1 < m/q < 10^{10}$ ) in a very high density nonneutral plasma (densities exceeding the Classical Brillouin Limit), and makes recommendations for future work.

The central problem is the "containerless" and "collisionless" assembly of subatomic, atomic, and molecular fragments into bulk antimatter, in this case solid antihydrogen "ice." It is necessary to condense antimatter into this form to achieve favorable payload fractions (when it is used as a fuel for propulsion) and to have a new means of high-density energy storage (when it is used for energy storage). Since this condensation has not been done before and must be done under "containerless" and "collisionless" conditions (with regard to collisions with ordinary matter), severe restrictions are placed on the methods used. Fortunately, normal matter can be used as a simulant in all aspects of the investigation, since the only relevant difference in the two forms of matter in this case is the sign of the atomic charge (ignoring parity and time). All work described in this report will refer to normal hydrogen, unless otherwise denoted.

As a result of our research for the past two years, an appropriate condensation method is now known whereby high density clusters may be grown via "containerless" condensation methods. This method, the CI approach, uses condensation of suitable reactants upon cold, trapped, hydrogen cluster ions.

To perform this "containerless" condensation (CC), cold fragments such as hydrogen atoms, molecules, and ions must be formed (resulting in the release of significant amounts of chemical energy) that have little or no translational, vibrational, or rotational energy. These cold fragments are then assembled to form ever larger fragments. The end result of this research will be a detailed and experimentally established "containerless" synthetic method.

# TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION	1
The Neutral Reactants: H and H <sub>2</sub>	3
CLUSTER ION APPROACH TO CONTAINERLESS CONDENSATION	7
Hydrogen Cluster Ions: Review of Structure and Properties	7
Large Hydrogen Cluster Ion Sizes and Trapping	9
Containerless Condensation by Pseudo-Wall Reactions	9
<u>Introduction to Pseudo-Walls</u>	9
<u>Condensation Pathways Using Pseudo-Walls</u>	14
Destruction of Hydrogen Cluster Ions by Annihilation	17
Very Cold Hydrogen Cluster Ions	22
EXPERIMENTAL DESIGN	25
Vacuum Chamber Components	26
Ion Trap Specifications	29
Pumping System	34
Transit Spreading of the Cluster Ion Beam	34
The Wien Filter	36
The Sector Magnet	37
Cluster Ion Detection	40
The Cluster Ion Source	42
System Performance	43
CONCLUSIONS AND RECOMMENDATIONS	51
REFERENCES	55
APPENDIX A	A-1
APPENDIX B	B-1

## LIST OF FIGURES

<u>Figure</u>	<u>Caption</u>	<u>Page</u>
1	Flowchart of Events that Led to the Proposed Plan of Research	2
2	Energy Levels of Atomic and Molecular Hydrogen	4
3	Reaction Matrix Showing Containerless Condensation Routes	13
4	Reaction Matrix with a Potential CC Pathway	15
5	Reaction Matrix with a Potential CC Pathway	16
6	Reaction Matrix with a Potential CC Pathway	17
7	Reaction Matrix with a Potential CC Pathway	18
8	Annihilation Cross-Section Data	20
9	Diagram of the Overall System Layout	25
10	Schematic of the Main System Components	26
11	Schematic Showing a Close View of the Source Chain	31
12	Diagram of the Ion Trap Showing the Trap Walls, End Caps, Grids, and Central Rod	32
13	Diagram of the Entire Trap Section Showing the End and Side Projections	33
14	Pumping System and Beam Spread Model	35
15	Diagram of the Wien Filter Showing Configuration and Construction Design	36
16	Diagram of the Sector Magnet Showing Configuration and Construction Design	38
17	Linear Plot of Magnetic Field Variation Along the Axis of the Superconducting Magnet (Normalized to the Central Field)	39
18	Log Plot of Magnetic Field Variation Along the Axis of the Superconducting Magnet (Normalized to the Central Field)	39
19	Schematic of the Detection System Intended for Use in the First Experiments	40

# LIST OF FIGURES (Con't.)

<u>Figure</u>	<u>Caption</u>	<u>Page</u>
20	Proposed Cluster Ion Sources	44
21	Cyclotron Frequency Versus Cluster Number	45
22	Central Rod Potential Versus Cluster Number at Constant Total Mass	46
23	Plasma Frequency Versus Cluster Number	47
24	Plasma Number Density Versus Cluster Number	49
25	Mass Density Versus Cluster Number	49
26	Total Mass Confined by the Trap Versus Cluster Number for a 100-cm Trap	50
27	Total Mass Confined by the Trap Versus Cluster Number for a 300-cm "Next-Generation" Trap	50
A-1	Plot of the Two Rotational Frequencies of a "Solid" Plasma and the Location of the Brillouin Limit	A-2



# LIST OF SYMBOLS

$2\alpha$	# nuclei in a cluster ion or mass of a cluster per proton mass
$\alpha$	# molecules in a cluster ion
$n$	number density $\text{cm}^{-3}$
$V$	volume
$v$	velocity
$N$	# of cluster ions
$M$	mass, critical size for $M^+$ process
$a$	radius of sector magnet, S.C. magnet radius, critical site $A^+$ , central conductor radius
$D$	$\text{cm}^3/\text{molecule}$ for hydrogen cluster ions
$z$	charge per cluster, axial distance from center of S.C. magnet
$z$	distance, spacing between electrodes in Wien filter
$L^0$	mean free path
$V'$	potential
$\beta$	charge state $\pm$
$f$	collision frequency
$L'$	attenuation coefficient
$e$	charge of electron
$c$	conductances (L/s)
$I$	current (amps)
$J$	flux
$A$	area, Einstein A
$B$	magnetic field
$E$	electric field, energy
$\phi$	potential on central rod

# LIST OF SYMBOLS (Con't)

$\Omega, \omega_c$	cyclotron frequency
$d$	radius of trap
$R^\pm$	annular plasma inside ( $R^-$ ) and outside ( $R^+$ ) radii
$k$	Boltzmann constant, integer
$T$	temperature
$\Gamma$	plasma state parameter
$r_{ci}$	cluster ion radius
$r_L$	Larmor radius
$q$	charge
$S_p$	speed of pump
$P_s$	source pressure
$P_t$	trap pressure
$\alpha_w, R_w$	outgas rate (watts/m <sup>2</sup> )
$A_w$	wall area
$w$	slit width
$R$	radius of curvature
$L$	length of S.C. magnet (axial)
$Q$	cross section, throughput of pumping section
$r$	radius of annular plasma
$p$	integer
$\Delta$	thickness of plasma (annular)
$\omega_p$	plasma frequency
$v_\theta^o$	azimuthal velocity
$E_r^o$	radial electric field
$u$	mass to charge ratio

## INTRODUCTION

The future development of high-power, high-flux accelerators is projected to result in the eventual production of significant annual yields (milligrams) of antiprotons and positrons. The availability of this new form of matter suggests a fascinating new science having many applications. One application of this new substance (in the area of energy storage) uses it as a propellant with unsurpassed characteristics (payload fractions, specific impulse, thrust etc.) which make it ideally suited for missions currently impossible. However, before it can be used in this application, it must be put into a high-density form; it must somehow be nucleated (or condensed) into a form that leads to high energy density and is easily confined (levitated). Fig. 1 is a diagram of the approach for the past two years. R. L. Forward (Refs. 1,2) has suggested frozen parahydrogen (levitated) as an ideal form; however the means by which one obtains this product (starting with trapped protons and electrons) was unknown. Hence, project research for the past two years sought a means of bridging the enormous gap between individual, unrecombined fermions and bulk matter. The CI approach to "Containerless Condensation" (CC) soon emerged as a feasible approach within constraints imposed by present-day and near-term technology. The CI approach uses the growth and manipulation of hydrogen cluster ions in traps using electric and magnetic fields. The main advantages of cluster ions are these:

1. They are more stable than uncharged clusters (except  $H_2^+$  versus  $H_2$ ). Cluster ions with mass to charge ratios beyond  $u=10^5$ , have been observed experimentally (Refs. 3,4).
2. They can be trapped, transported, and stored in devices that use the Coulomb force (trap times of months and depths greater than 10 eV are typical) (Refs. 5,6).
3. They have very large rate constants for collision with a neutral antiparticle ( $>10^{-9} \text{ cm}^3 \text{ s}^{-1} \text{ particle}^{-1}$ ).
4. They have a dipole-allowed IR spectrum and thus may radiatively cool their vibrational degrees of freedom (except  $H_2$ ) (Einstein A's  $\sim 10^3$




Figure 1. Flowchart of Events that Led to the Proposed Plan of Research.  
Also given are the names associated with each category.

$s^{-1}$ ). The spectroscopy of normal  $H_3^+$  (the analog of  $H_3$ ) has been reported (Refs. 7-10). The radiative properties have been reviewed, (Refs. 11-13) as have large cluster ions (Ref. 14).

5. Their translational degrees of freedom may be cooled by existing (nonlaser) techniques.

The HCISG (Hydrogen Cluster Ion Study Group) meetings (Ref. 15) reinforced enough of the original notions to deserve a workshop (called CCSHCI, for Cooling Condensation and Storage of Hydrogen Cluster Ions) devoted to a more in-depth look. Three general strategies emerged: radiative, nonradiative, and both. For the most part, there seem to be no promising stimulated radiative channels (and few spontaneous channels), due primarily to the general lack of bound excited electronic states in these systems, as shown in recent calculations on  $H_3^+$  (Ref. 16). Hence, one is led to consider nonradiative pathways with the hope that spontaneous pathways will take care of themselves. The pseudo-wall concept devised by the author and Professor W. C. Stwalley has been implemented (in collaboration with Professors A. Y. Wong and G. Rosenthal) as discussed in the Experimental Design section.

The Neutral Reactants: H and  $H_2$

A brief review of these molecules in the context of CC is presented. How to proceed with the radiative formation of the simplest antifrags (H,  $H_2^+$ ,  $H^-$ ,  $H_2$ ) is still rather uncertain. For  $H_2^+$  formation see Helm's report (Ref. 17). Since  $H^-$  does not have discrete excited states, no way is known to manipulate it radiatively. The formation of  $H^-$  may work well in a pseudo-wall environment. The small negative cluster ions have been removed from consideration due primarily to their instability (Refs. 13, 18).

Hydrogen atoms can be formed via stimulated recombination (the most direct path being continuum-2p) followed by spontaneous radiation (2p-1s) to the ground state (Fig. 2) (Refs. 19, 20). Hydrogen atoms may also be laser cooled to near the quantum limit (Ref. 21), and otherwise laser

size were available, one might use them as a substrate for the recombination of hydrogen atoms. Ideally, one would like to nucleate hydrogen atoms directly onto (or into) a cluster ion. Research is needed that reveals how the recombination energy is coupled into the cluster ion. Once formed, however, the means of trapping the atoms are known (Refs. 23-25).

Unless a breakthrough occurs, any nucleation process that uses hydrogen molecules is likely to be complicated and inefficient. A specific nucleation pathway that uses H and avoids  $H_2$  is actively being sought. For  $H_2$  to be useful in the nucleation of the smallest hydrogen cluster ions (say,  $u < 29$ ), it must be quite cold in all of its degrees of freedom, which is difficult to accomplish. Further difficulties are that  $H_2$  cannot radiatively cool itself in its ground state (without the help of a third body) and we know of no simple way of laser cooling any of its degrees of freedom.  $H_2$  formation is difficult and inefficient and once formed there is no known way of keeping it from annihilating on the walls (see report by K. M. Sando, in CSHCI proceedings). The means of making it radiatively are currently being studied (Ref. 26). For a discussion of the intramolecular potential, see Lester (Ref. 27).

## CLUSTER ION APPROACH TO CONTAINERLESS CONDENSATION

This section is devoted to topics that relate in one way or another to the development of CC techniques.

### Hydrogen Cluster Ions: Review of Structure and Properties

As stated in the Introduction, the properties and structure are well known for the smallest cluster ions and have been reviewed extensively (Stwalley (Ref. 28) has prepared an extensive bibliography; see Appendix B). Some relevant properties of cluster ions are included in Table 1 (Ref. 29).

As a review, we are considering just the positive cluster ions. This was decided because the small negative cluster ions (excluding  $H^-$ ) are unstable to spontaneous detachment. Although it is known that negative hydrogen cluster ions become stable at some (unknown) critical size, they would still be sensitive to photodetachment, making them less desirable (Ref. 13).

In current penning-type trapping systems a negative ion plasma is more difficult to deal with. With a positive ion plasma, a small electron population will simply move to the ends of the trap and be removed by the end termination electrodes. With a negative ion plasma, a small population of electrons would not be able to leave the system. These trapped electrons add charge to the system and introduce the possibility of two species diocotron instability. Such instability would probably reduce confinement time.

The small singly charged, positive hydrogen cluster ions are well studied both experimentally and theoretically, and are all known to be stable. The doubly charged hydrogen cluster ions are stable for sizes larger than about 2000 (this may be inferred from data taken from Yano [Ref. 4]). Calculations predict the dipositive ions to be stable at size 887, tripositive ions at 2906, and the tetrapositive ions at 6196

TABLE 1. Overview of Particle (and Antiparticle) Properties and Characteristics. (From Stwalley [Ref. 29]).

particle	$e^-$	$H^+ (p)$	$H$	$H^-$	$H_2^+$	$H_2$	$H_3^+$	$H_3$	$H_4^+$	$H_4$	$H_{N+}^+$	$H_{N+}$
antiparticle	$e^-$	$H^+ (p)$	$H$	$H^-$	$H_2^+$	$H_2$	$H_3^+$	$H_3$	$H_4^+$	$H_4$	$H_{N+}^+$	$H_{N+}$
binding energy in eV	-	-	13.6	0.75	2.6	4.5	4.1	0	0.1	<0.01	...0.008	0.008
products	-	-	$H^+, e^-$	$H, e^-$	$H^+, H$	$H, H$	$H^+, H_2$	$H, H_2$	$H, H_3$	$H_2, H_2$	$H_{N-2}^+, H_2$	$H_{N-2}, H_2$
sensitivities <sup>†</sup> :												
$\lambda < 0.2 \mu$ VUV	-	-	x	x	x	x	x	x	x	x	x	x
0.2-0.4 $\mu$ UV-VIS	-	-	-	x	x	-	(x)	-	(x)	-	x	x
0.7-100 $\mu$ IR	-	-	-	-	-	-	-	-	-	-	x	x
$\lambda > 100 \mu$ microwave	-	-	-	-	-	-	-	-	-	-	x	x
radiative recomb. ( $e^-$ )	*	x	x	-	(x)	-	(x)	-	(x)	-	(x)	-
dissoc. recomb. ( $e^-$ )	*	-	-	-	x	-	x	-	x	-	x	-
assoc. detach. ( $H$ )	-	-	x	*	-	-	-	x	-	-	-	-
rxn. ( $H$ )	-	-	*	-	x	-	x	(x)	x	-	-	-
rxn. ( $H_2$ )	-	-	-	-	x	*	-	-	x	-	-	-
neutralization ( $H^-$ )	-	x	-	*	x	-	x	-	x	-	x	-
radiative assoc. ( $H, H_2$ )	-	-	-	-	-	-	-	-	-	-	x	x
trapping:												
Penning/Paul laser	x	x	-	x	x	-	x	-	x	-	x	-
inhomogeneous magnet	-	-	x	(x?)	x?	x?	-	-	-	-	x	x
cooling:												
radiative laser	-	-	-	-	-	-	x	-	x	-	x	x
lepton ( $e^-$ )	x	x	-	x	-	-	-	-	-	-	?	-
transfer:												
electrostatic	x	x	-	x	x	-	x	-	x	-	x	-
magnetic	-	-	x	-	-	-	-	-	-	-	-	x
laser	-	-	x	-	(x?)	(x?)	-	-	-	-	-	-

\* N assumed large ( $>10^3$ ).

† All collision partners assumed to be cold.

\* Reactant.

Refs. 30, 31). The odd-numbered cluster ions consist of an  $H_3^+$  core, surrounded by  $H_2$  molecules. The even-numbered cluster ions are similarly structured but have one attached H atom.

#### Large Hydrogen Cluster Ion Sizes and Trapping

Taking the density of solid hydrogen ( $0.076 \text{ g cm}^{-3}$ ) to represent the density of a large cluster ion, one obtains the constant  $D = 5 \times 10^{-23}$



cm<sup>3</sup>/molecule, which was used to construct Table 2. In principle, if the correct source of hydrogen cluster ions is developed, virtually any sized cluster ion may be trapped in the proposed ion trap, up to roughly

TABLE 2. Large Hydrogen Cluster Ion Diameters.

<u>Cluster #</u> <u>(# of molecules)</u>	<u>Log (cluster diameter)</u> <u>(microns)</u>
10 <sup>4</sup>	-2
10 <sup>7</sup>	-1
10 <sup>10</sup>	0
10 <sup>13</sup>	1
10 <sup>16</sup>	2
10 <sup>19</sup>	3

10<sup>14</sup> atoms. This is discussed further in the section entitled "Very Cold Hydrogen Cluster Ions."

#### Containerless Condensation by Pseudo-Wall Reactions

The ideal way to perform containerless condensation is to arrange conditions whereby both radiative and nonradiative processes occur. This section briefly discusses the concept of a pseudo-wall and presents a number of condensation pathways based on this concept.

#### Introduction to Pseudo-Walls

The primary goal is a detailed understanding of the available association channels of hydrogen cluster ions. The simultaneous requirements of the association reactions are the release of energy,

conservation of momentum, and generation of a larger cluster ion. Generally, the association pathways may be categorized as radiative, for example,



or nonradiative,



Both of these reactions are exothermic. However, in reaction (1a), a larger cluster ion is obtained because the channel allows the excess energy (kinetic, electronic, etc.) to be radiated as a photon. In case (1b), the excess energy is released in the form of kinetic energy of the ejected atom. Certain channels may exist that involve both kinds of processes simultaneously. These channels are still being explored.

One way of ensuring that collisional and radiative relaxation channels are favored is to manipulate the reactant densities and temperatures, i.e., have a reaction that always involves one (or both) of the reactants in a special condensed phase, referred to as a "pseudo-wall." A pseudo-wall is a cold, trapped, high-density phase that is thermally connected (via E and M fields) to the environment. This cold, high-density condensed phase results in collective effects that may occur on short time scales; it behaves somewhat like a wall both kinetically and thermodynamically. This concept is really just an extension of the liquid and solid pure electron plasmas that have been talked about for some time (Refs. 32, 33). The relevant parameter is

$$\Gamma = e^2 / (a kT),$$

where  $a$  is given by  $[(\frac{4}{3}) \pi a^3 n] = 1$ ,  $e$  is the charge,  $T$  is the temperature,  $n$  is the density, and  $k$  is Boltzmann's constant.  $\Gamma=1$  defines a plasma, at  $\Gamma=2$  one has a liquid, and finally at  $\Gamma=155$  one obtains a solid pure electron plasma.  $\Gamma$  is a measure of the Coulomb potential to the

thermal kinetic energy in the plasma. One wants the largest range of  $\Gamma$  obtainable. This may be done by varying  $T$  and  $n$ .

For the proposed device, the achievable density will be roughly  $n=10^8 \text{ cm}^{-3}$  for clusters smaller than  $10^4$  atoms. If the  $T=4 \text{ K}$ , one obtains

$$\Gamma \sim 120,$$

which falls well within the liquid regime.

When a single incoming (neutral or charged) particle collides with the pseudo-wall, part of the collision energy is absorbed, almost instantaneously, by neighboring particles that make up the wall. The other important feature is that, since the pseudo-wall is connected to an ion trap by  $E \& M$  forces, it can transfer its energy to the environment. When referring to the above reactions, if just  $\text{H}_2^+$  is made into a pseudo-wall, they become



and



Here, we consider the reaction of  $\text{H}_2$  molecules within the cold, high density environment of  $\{\text{H}_2^+\}$  ( $T < 4 \text{ K}$  and  $n = 10^{11} \text{ cm}^{-3}$ ). The importance of this process is that the collisions are not strictly 2-body, but may be 3-body (and higher, up to  $10^{14}$  body for the proposed pseudo-wall). This results in substantial collisional energy transfer (rather than just radiative) of the energy of the exothermic reaction to the pseudo-wall (and ultimately to the environment). The advantage illustrated in this example is that reaction (1a) cannot proceed radiatively, while (2a) can. In this manner, a finite set of reaction pathways result that may proceed much more efficiently and at much higher rates (coming from the inclusion of collisional relaxation mechanisms, and high effective densities).

The conditions that specify a pseudo-wall are now defined. The main constraint is  $(L \ll r)$ , where  $L^0$  is the collisional mean free path of the cluster ion and  $a$  (assume  $a=1$  cm) is the physical radius of the pseudo-wall. If  $L^0 = .01a = 10^{-2}$  cm, and the collisional cross-sections ( $Q$ ) are at least  $10^{14}$  cm<sup>2</sup>, the pseudo-wall density ( $n$ ) is obtained from

$$L^0 = .001a = 1/(nQ),$$

giving a neutral density  $n = 10^{16}$  cm<sup>-3</sup> (the temperature is adjusted to achieve this density). In contrast, the maximum densities of ions in non-neutral plasmas are  $10^{11}$  cm<sup>-3</sup>. Under these conditions a single cluster ion in a pseudo-wall would undergo roughly  $10^6$  bimolecular collisions per second with neutral particles. Constructing such a cylindrically shaped pseudo-wall of volume ( $V=30$  cm<sup>3</sup>), would require only about  $N=nV$ , or  $3 \times 10^{12}$  cluster ions.

Another way of specifying pseudo-wall conditions is to require that the collision frequency,  $f$ , be greater than the radiative lifetime,  $t$ , of any excited states, ensuring collisional deactivation at some velocity  $v$ :  $f = nQv > 1/t$ . This criterion is satisfied for neutral densities,  $n$ , of  $10^{16}$  cm<sup>-3</sup>, cross sections,  $Q$ , of  $10^{-14}$  cm<sup>2</sup>, and velocities of  $10^4$  cm s<sup>-1</sup> for radiative lifetimes of  $10^{-6}$  s and longer (cluster ions have radiative lifetimes of several milliseconds, for vibrational transitions).

The maximum cooling power required of the pseudo-wall may be estimated. The most exothermic reaction is the first one in Fig. 3, requiring 13.6 eV for each H atom. Assuming a nucleation rate of  $10^{12}$  particles per second (166 nanoamps), this results in  $13.6 \times 10^{12}$  eV s<sup>-1</sup> (or 2 micro-watts) of recombination energy that must be coupled through the pseudo-wall to the environment. Established resistive cooling techniques will be used, if necessary.<sup>34</sup>

The main reactions which may take place in pseudo-walls are illustrated in the reaction matrix of Fig. 3. Certain working assumptions were made in the construction of this matrix:




Figure 3. Reaction Matrix Showing Containerless Condensation Routes.  
In the first reaction, protons are added to a pseudo-wall composed of electrons, resulting in the production of cold hydrogen atoms.

1. All particles (one or both reactants) can be turned into pseudo-walls; ionic pseudo-walls are currently preferred.
2. Only two-reactant combinations are considered (reactant plus pseudo-wall).



3. No nuclei (other than deuterium) larger than the proton.
4. Minimum dependence on radiative processes (photons are excluded).
5. No anions other than  $H^-$ .
6. No obvious fragmentation reactions.
7. No unstable or short-lived particles (for example positronium,  $H_2^-$ ,  $H_3^-$ , etc.).

Some reactions in Fig. 3 (with asterisk) could proceed spontaneously within the same pseudo-wall to grow large cluster ions. Recall that the cluster ions of sizes  $H_m^+$  and  $H_a^+$  are the threshold cluster ion sizes for ion-molecule ( $M^+$ ) and ion atom ( $A^+$ ) association, respectively. These have been postulated to exist at sizes around 29 and 2000 respectively.<sup>15</sup> The objective is to select the optimum pathway to these large cluster ions by connecting reactions and defining pseudo-walls.

#### Condensation Pathways Using Pseudo-Walls

Fig. 4 illustrates one of the many possible condensation pathways. One example (out of many) of a "containerless" condensation pathway is designated by the large arrows in Fig. 4. The proposed experimental approach is designed to verify the reactions and pathways of this matrix. The most fundamental reactions are those involving either  $H^+$  or  $e^-$  which react to produce  $H$ ,  $H^-$ , or  $H_2^+$  (the production of  $H_2$  may also occur in these cold plasmas but this is assumed minimal). The reactions indicated by the condensation pathway in Fig. 4 may be written specifically as

1.  $(e^-) + H^+ = H$
2.  $(e^-) + H = H^-$
3.  $(H^-) + H = H_2 + e^-$



The procedure is repeated until a critically sized cluster ion is obtained such that it will undergo repeated additions of  $H_2$  without fragmentation. This may be written






Figure 4. Reaction Matrix with a Potential CC Pathway. (Note large arrows.) The reaction denoted by ( ) may be written,  $H^+ + p H_2 = H_{p+1}^+$ , where  $p$  is an integer. Refer to text for reactions denoted with numbers.

$H_m^+$  is referred to as the threshold seed cluster ion size for the  $m^+$  process. If this process is repeated enough times another seed critical cluster ion size will be reached such that the reaction



will proceed. Experiments have been proposed that will determine the cluster ion sizes  $H_a^+$  and  $H_m^+$ . A few other possible pathways have been illustrated in Figs. 5 to 7.




Figure 5. Reaction Matrix with a Potential CC Pathway.  
(Note large arrows.)






Figure 6. Reaction Matrix with a Potential CC Pathway.  
(Note large arrows.)

#### Destruction of Hydrogen Cluster Ions by Annihilation

This section discusses the various ways hydrogen cluster ions may be destroyed or removed from the trap due to collisions (resulting in surface annihilation) with neutral (and nonneutral) antihydrogenic species (made up of fermions, these represent a main constituent of background gas).







Figure 7. Reaction Matrix with a Potential CC Pathway.  
(Note large arrows.)

Considering just the primary event, the only annihilation products to consider are high-energy pions (Refs. 1, 2) from the proton-antiproton annihilation and  $\Gamma$  rays from electron-positron annihilation. It is assumed that the coupling of motion between Pions and the molecular (and atomic) degrees of freedom is small, especially vibration, rotation, and translation (sources of pion-electron cross sections for electronic excitation, ionization, should be found). The  $1/e$  attenuation coefficient for  $\gamma$  rays is much larger ( $L'(1/e) > 128$  cm,  $E > 0.5$  MeV) than the dimensions of the cluster ions or plasmas and may be neglected.

There are other ways of destroying cluster ions: recombination, photodissociation, etc.; these however have been considered elsewhere (Refs. 9, 35). The reactions to be considered here are



+ annihilation products, and



Reaction 1 may be neglected for low-energy positrons.

Reaction 2 under the above assumptions will not yield fragments, but will only reduce the mass of the cluster ion by one unit; it too is neglected.

Reaction 3 is the most problematic since it leads to the production of a neutral cluster ion,  $\overline{H_{n-1}^0}$  (only slightly less massive). Here, we must consider the cross section for proton-antiproton annihilation. Considering destruction by just low energy (thermal  $T=300$  K) antiprotons, an extrapolation of the data (Ref. 36) from Fig. 8 (upper right) gives a cross section of  $10^9$  barns or  $10^{-15} \text{ cm}^2$ . The density of protons in a cluster ion is roughly  $10^{22} \text{ cm}^{-3}$  so that the mean free path of an antiproton in a cluster ion can be estimated to be

$$L = 1/(10^{-15} \times 10^{22}) = 10^{-7} \text{ cm}.$$

This means one need only consider surface annihilation from thermal antiprotons since the cluster ions will, in general, be much larger than 1 nm. Since the populations and annihilation cross sections for higher




Figure 8. Annihilation Cross-Section Data. (Figure courtesy of V. Flaminio, W. G. Moorhead, D.R.O. Morrison, and N. Rivoure, CERN).

energies decrease rapidly (with increasing energy), they may be disregarded.

The fate of the particle produced by reaction 3,  $H_{n-1}^0$ , is now considered in the low-temperature, high-magnetic-field environment of the cylindrical "Penning" trap. The resulting cold neutral cluster,  $H_{n-1}^0$ , may be either even or odd, and have different ortho to para ( $H_2$ ) ratio. The possible categories are summarized:

<u>Case</u>	<u>Even</u>	<u>Odd</u>	<u>Para</u>	<u>Ortho</u>
1	x		x	
2	x			x
3		x	x	
4		x		x

In cases 1 and 2, the cluster has a single unpaired electron, giving it an appreciable magnetic moment (1 Bohr magneton). The neutral cluster would behave like a very large hydrogen atom and hence all the magnetic schemes developed for H atoms would also apply. Although the cluster would be confined by a shallow axial barrier (near the ends of the magnet), it would drift radially until it encountered something. It is interesting that, if by chance it encountered the pseudo-wall before reaching the ion trap wall, it should be recaptured by the pseudo-wall.

In cases 3 and 4, the cluster magnetic properties are three orders of magnitude weaker than in cases 1 and 2, being due only to the coupling of the nuclear spins (1 nuclear magneton). In case 3, the cluster is composed mainly of parahydrogen; it is diamagnetic (negative susceptibility), has only even rotational levels, and is a "low field" seeker. The magnetic field is essentially constant in the radial direction so the cluster will tend to be gently expelled axially out of the trap. Since the trap is essentially solenoidal, there will be no appreciable radial force. Once more, it may either encounter the pseudo-wall first, or it may just go out of the trap. If the cluster leaves the trap, it could conceivably be recaptured following laser photoionization. In case 4, the resulting cluster is mostly Orthohydrogen; it is paramagnetic,

(positive susceptibility), has only odd rotational levels, and is a "high field" seeker. This means that there will be a gentle restoring force tending to keep the cluster in the high-field region of the trap. If the initial kinetic energy of the cluster is not greater than this magnetic barrier, the cluster would orbit about the trap center until it encountered the pseudo-wall and recondensed, or hit the trap wall.

The destruction of cluster ions by background hydrogenic species has been found to be something of concern, but does not imply insurmountable difficulties in the pseudo-wall concept.

#### Very Cold Hydrogen Cluster Ions

If cluster ions of any size may be trapped, one might well ask: Under what circumstances is the Larmor radius smaller than the physical radius of the cluster ion? In this situation the internal motion of the charge on the cluster ion is coupled to the (now very low frequency) cyclotron motion. The remaining degrees of freedom are just a slow  $\vec{E} \times \vec{B}$  drift plus axial motion. The whole plasma should be quite dense and fulfill some of the main requirements of a pseudo-wall.

The radius of a spherical cluster ion is

$$r_{ci} = \{(3V)/(4\pi)\}^{1/3} = \{(3D\alpha)/(4\pi)\}^{1/3},$$

where  $D$  is  $5 \times 10^{-23} \text{ cm}^3/\text{molecule}$  and  $\alpha$  is the number of molecules in the cluster. The constraint is that  $r_{ci}$  be less than the Larmor radius ( $r_L = (2\alpha m v)/(qB)$ ), or

$$r_L < r_{ci},$$

where  $(2\alpha)$  is the cluster ion size,  $m$  is the mass of a hydrogen dimer,  $v$  is the perpendicular component of the cluster ion velocity (includes radial and azimuthal motion),  $q$  is the charge, and  $B$  is the magnetic field.

This leads to the following inequality;

$$(2\alpha m v)/(qB) < 2.3 \times 10^{-6} (2\alpha)^{1/3} \text{ (meters).}$$

Putting in values for m, q, and B (8 Tesla), gives

$$[(3.3 \times 10^{-27})/(2.3 \times 10^{-6} \cdot 1.6 \times 10^{-19} \cdot 8)] (2\alpha n^{2/3} v) < 1.$$

Simplification then yields

$$(2\alpha)^{2/3} v < 890 \text{ (m/s),}$$

where v is in (m/s). This result (which is increasingly less accurate for small  $\alpha$ ), is given in Table 3 below (with a column of temperatures, assuming a thermal source).

Since the proposed trap will have a liquid helium baffle for cooling, it should be capable of producing this cold, slowly rotating pseudo-wall for cluster sizes  $\alpha < 5000$  dimers (or cluster sizes,  $2\alpha < 10^4$ ).

TABLE 3. Velocities and Temperatures of Cluster Ions  
with  $r_L < r_{ci}$ .

<u>Log(n) (dimers)</u>	<u>v (m/s)</u>	<u>T (K)</u>
1	192.	34.6
2	41.4	16.0
3	8.92	7.46
4	1.92	3.46
5	0.414	1.61
6	0.0892	0.750
7	0.0192	0.346
8	0.00414	0.160

## EXPERIMENTAL DESIGN

The system described is similar to the one proposed recently by Kluge et al (Ref. 37). However, since it will contain an annular plasma, it has many of the design features developed by Wong and Rosenthal (Ref. 38).

A generalized schematic of the system is shown in Fig. 9. Note that the main components of the vacuum chamber are in an "L" configuration. It is divided (top to bottom) into the diagnostics, ion trap, 6-way cross, and source chain. To the sides of the 6-way cross are attached the Combo Vac, Cryo Pump (optional), and Star Flange.




Figure 9. Diagram of the Overall System Layout. The main system is divided into main sections; the source chain constitutes the entire lower section beneath the 6-way cross.



Attached to the bottom is a long, narrow, straight section that makes up the source chain. Note that many of the system components must be bakable (B) to 700 K. The remaining components are the ion gauge and N<sub>2</sub> source. The system is designed to operate satisfactorily with just these mentioned components; however, the optional system, using a mechanical/sorption pump combination is recommended as a later "add on." The addition of the optional system is recommended and would lengthen the times between servicing the Combo and Cryo pumps.

#### Vacuum Chamber Components

The main components of the vacuum chamber are shown in Fig. 10.




Figure 10. Schematic of the Main System Components. (See text.)

There are 16 essential parts. Unless otherwise noted, 304 stainless steel is to be used for these components, most of which are bakable (excluding pumps) (see Fig. 9). The parts are discussed separately with regard to function and size. This information is from known commercial sources; part numbers were taken from Thermionics Vacuum Products, Hayward, CA; Norcal is suggested as a competitor.

1. The trap wall section is custom built (Fig. 10). It is bakable and features a slip joint lower end (item 7) to allow assembly through the bore of the superconducting magnet (that is, the lower rotatable flange is removable, split into two halves, so that the trap wall section can pass through the magnet bore). This feature will allow easy disassembly. The top flange is a standard rotatable 8 in. conflat.

2. Standard 6-way cross. This component joins the ion trap, combo vac, star flange, optional cryopump, and source chain. One port is unused for access (or future addition).

3. Combo-Vac pump. Commercially available, it comes equipped with a hydrogen diode pump, an 8 in. OD flange, and should be rated at either 750 L/s or 1000 L/s.

3a. 10 in. to 8 in. adaptors will be needed if either the 1000 L/s Combo pump or Cryo-Torr 10 pumps are used.

3b. Bakeout heaters (4) for the 6-way cross. These include controllers.

3c. Liquid nitrogen level controller.

4. Cryopump. The optional cryopump may be added to the available port (suggest CTI Cryo-Torr 10). Space is reserved for this pump, should better vacuum or more gas load (than expected) result.

5. Star Flange. This is commercially available and may be custom built. It provides four smaller ports that will be used for liquid

helium feedthrough (for the liquid helium baffle), controls for the ion trap, N<sub>2</sub> feedthroughs for flooding the system with 1.0 atmospheres of nitrogen, and control of the quadrupole deflector (item 5). This is a 10 in. conflat to 2 3/4 in. (x4) conflat.

5a. One blank 2 3/4 in. conflat.

5b. Nude ionization gauge for monitoring trap pressure. This should have Iridium filaments.

5c Flange for mounting the liquid Helium feeds for liquid helium baffle coolant.

5d. Bakable valve with 2 3/4 in. conflat. Flange with bakable air valve connects to N<sub>2</sub> with safety valve for air vent.

6. Quadrupole lens. This consists of four plates in quadrupolar arrangement and may be home built. Passing the cluster ion beam through this lens causes the beam to take on a helical trajectory. This is used for injecting an annular plasma into the trap.

7. Conflat flange. 10 in. to 4 1/2 in. adaptor.

8. Nipple with 4 1/2 in. conflat.

9. Tunable, Electromagnetic Sector Magnet. This should be custom built by a vendor (suggest Wang NMR) (see also the section on sector magnets). The maximum magnetic field is 0.5 T.

10. Nipple with 4 1/2 in. conflat.

11. Air driven, bakable gate valve with 4 1/2 in. conflat.

12. Nipple with 4 1/2 in. conflat.

13. Tunable Wien filter with electromagnet. This features a yoke for confinement of its magnetic field and should be custom made (suggest Wang NMR) (see also the section on Wien filters).

14. 4-Way Cross with 4 1/2 in. conflat flanges.

15. Nude ionization gauge for monitoring source pressure. This should have Iridium filaments.

16. Hydrogen cluster ion source. An appropriate source that will work is not commercially available and will have to be developed (see section on cluster ion sources).

Table 4 lists the specifications and prices for many of the items.

Fig. 11 shows the locations of the slit assemblies for the Wien filter and the sector magnet. Also given are the specifications for the magnets and electrodes of the Wien filter. Notice that an empty flange is attached to the sector magnet; this is reserved for the addition of a cryopump should the source produce excessive gas load on the pump(s) attached to the 6-way cross.

Also shown is the bakable gate valve used for isolating the trap environment from the source for long-term studies of trapped cluster ions in ultra-high vacuum and the sector magnet.


#### **Ion Trap Specifications**

The cylindrical ion trap is a modified version of devices used at UCLA and at UCSD. It is designed specifically for large mass-to-charge ratio cluster ions and features a central conducting rod (Fig. 12) that will be used for the neutralization of space charge in the confinement of high-density plasmas. The copper conducting rod (0.6 cm diameter) is supported only at the top, and passes through the end cap grids (end termination grids are used to provide stability for the annular plasma);

TABLE 4. Specifications and Costs of Parts.

<u>Item</u>	<u>Size/Part Number</u>	<u>Unit Price</u>
1	OD 8 in., length 16 in.	\$ 900
2	OD 8 in.	3150
3	T#COV750, 750 L/s	8300
	T#COV1000, 1000 L/s	9500
3a	T#1000x800	270
3b	T#IH-100	75
3c	T#LNC-400	475
4		
5	10 in. conflat to four 2 3/4 in.	
5a	2 3/4 in.	14
5b	T#NG-1001	270
5c	Custom	
5d	T#BVV-152	795
5e	T#F275000V	230
6	Custom	
7	T#1000x450	245
8	4 1/2 in.	185
9	Custom	
10	4 1/2 in.	185
11	4 1/2 in. (custom)	1400
12	4 1/2 in.	185
13	Custom	
14	4 1/2 in.	350
15	T#NG-1001	270
16	Custom	

Important note: Due to the the welding and construction techniques used in Ultra High Vacuum systems, this system should never be subjected to more than 1 atmosphere of pressure; doing so could cause leaking.



1. Magnets: Length=20 cm, width=4 cm, spacing=2 cm,  $B=0.4$  T  
Electrodes: Length=20 cm, width=1.2 cm, spacing=5 cm
2. First entrance slit: 5.5 cm from the input electrode edge  
(1.0 mm x 6.0 mm)
3. Second entrance slit: 1 cm from input electrode edge (0.3 cm x 6.0 cm)
4. Exit slit: 1 cm from output electrode edge (0.3 cm x 6.0 mm)
5. Sector magnet slits: (1.0 mm x 6.0 mm)

Figure 11. Schematic Showing a Close View of the Source Chain.  
Also given are the dimensions of the Wien filter and the five slit assemblies.

it is not electrically connected to the end caps. Notice also that the rod support is hollow so that ions may drift, unencumbered, through the entire trap. Surrounding the central conductor (Fig. 13) is the annular plasma, followed by the inside wall of the trap which is 4 in. in diameter.

The trap electrodes are composed of polished, gold-plated copper (oxygen free) (thickness, 0.7 cm) and are electrically insulated yet thermally connected, via ceramic spacers) to a liquid helium baffle which




Figure 12. Diagram of the Ion Trap Showing the Trap Walls, End Caps, Grids, and Central Rod.

surrounds it. This was done to eliminate the effects of black-body radiation on stored cluster ions. There is a 1/2 in. gap (1/2 inch in radius) between the trap OD and the inside wall of the liquid helium baffle. The 1/2 in. gap is to prevent heat load on the liquid helium baffle from external sources. This space can also be used for connectors to the diagnostics. The region surrounding the baffle can be filled with porous carbon to improve system pumping of He. The liquid helium baffle is supported by ceramic spacers to the chamber wall (8 in. OD 304 stainless steel). The ID (bore) of the superconducting magnet is roughly 9 in.




Figure 13. Diagram of the Entire Trap Section Showing the End and Side Projections.

(length 20 cm), allowing room for the slip joint assembly of the lower flange during disassembly of the ion trap wall section. The Superconducting magnet should be custom built to specifications (suggest Oxford). The proposed magnet will use Niobium Titanium for generating fields up to 8.5 Tesla, with long-term use at 7.0 Tesla and with a stability of 3 parts per billion per hour drift. The magnet requires liquid nitrogen and liquid



helium as cryogens. The liquid nitrogen typically requires refill every 7 days (<200 cc/hr), and the liquid helium every 3-6 months (<15 cc/hr). At the outermost diameter are the Helmholtz coils, which may be fabricated at AFAL, and are used to make the magnetic field parallel with the trap axis.

#### Pumping System

The system pumping requirements were derived through the model shown in Fig. 14a (Ref. 39).  $P_s$  ( $10^{-7}$  Pa) is the assumed pressure at the cluster ion source, and  $C_1$  is the conductance of the source chain. This is deliberately made very small with the use of small apertures and long length.  $P_t$  is the pressure in the trap ( $10^{-9}$  Pa).  $C_2$  is the conductance of the channel connecting the trap to the source pump; this is made very large by using a large aperture and short length.  $P_p$  and  $S_p$  are the pressure and pumping speed of the source pump. The derived relation for  $S_p$  is

$$S_p = [(R_w A_w + C_1 P_s) - C_1 P_t] / P_t,$$

where the wall outgassing rate  $R_w$ , for baked stainless steel, is  $10^{-13}$  W/m<sup>2</sup>, and the inside area  $A_w$  of the system is  $1.26 \times 10^4$  cm<sup>2</sup>. Putting in the calculated values for  $C_1$ ,  $P_t$ , and  $P_s$ , gives  $S_p = 5340$  L/s (note that this pumping speed is exceeded in the pump specifications). This was done to ensure that the system will be easily able to achieve and maintain pressures in the trap below  $10^{-9}$  Pa. The derived relation for calculating the pressure in the trap is

$$P_t = (C_2 + S_p) (R_w A_w + C_1 P_s) / [(C_1 C_2 + S_p (C_1 + C_2))].$$

The derivation and details may be found in the Appendix.

#### Transit Spreading of the Cluster Ion Beam

In transit from the cluster ion source to the ion trap, the cluster ion beam will undergo a transverse spreading due to space charge

repulsion (Fig. 14b). The maximum allowable transit time of the cluster ion beam pulse is

$$t = 1.52 \times 10^{-3} [(R_f - R_i)/R_i] (1/z) (2\alpha/n)^{1/2},$$

where the maximum final radius is  $R_f$  (aperture of the ion trap), and the initial beam radius is  $R_i$  (aperture of the source), the charge per cluster is  $z$ ,  $2\alpha$  is the mass number of the cluster (mass of the cluster per proton mass), and  $n$  is the density of the cluster ions at the source. The constant has units of  $\text{sec cm}^{3/2}$  so that the answer is in seconds. With this relation, and knowing the distance to the trap, the mass of the transmitted ions, and the pressure of the source, one can easily calculate the acceleration potential ( $\Delta V'$ ) needed for the prescribed transit time. The derivation and further details are given in the Appendix.




Figure 14. Pumping System and Beam Spread Model. (a) Schematic of the overall system used for the pumping system calculation; (b) Schematic of the model used in the beam spread calculation (See Appendix).

## The Wien Filter

The Wien filter is a velocity filter that may use either a fixed or variable magnetic field (the latter field is preferred as an added degree of tuning flexibility). A schematic of a fixed field version<sup>4</sup> of the Wien filter is shown in Fig. 15. This figure is mainly to show device configuration and construction. The dimensions are given in the section entitled "Vacuum Chamber Components." Note that an iron field yoke is used to keep the magnetic field well confined so as to not perturb other system components. Also, not shown, it may be necessary to shield the Wien filter (and sector magnet) from the superconducting magnet by wrapping it with several layers of mu metal. In the construction of the Wien filter, a minimum number of feedthroughs and ceramic standoff insulators are used,




Figure 15. Diagram of the Wien Filter Showing Configuration and Construction Design.

since these have much larger outgassing rates than stainless steel (even when baked). The velocity of the ions transmitted through the filter is given by

$$v = 10^8 V'_d / zB$$

where  $V'_d$  is the potential between the electrodes with spacing  $z$ , and  $B$  is the magnetic field. The mass resolution of the Wien filter is given by (Ref. 40)

$$(\Delta m)/m = (2 z V' w) / V'_d L^2$$

where  $V'$  is the potential through which the injected ions have fallen,  $w$  is the width of the slits, and  $L$  is the length of the electrodes. A mass resolution of 0.01 is deemed sufficient.

#### The Sector Magnet

The electromagnetic sector magnet is a momentum filter which, when used in combination with the Wien filter, provides an output cluster beam that is both mass and velocity analyzed. A schematic of the device to be built is shown in Fig. 16. Shown are the input, output flanges (and the unused optional flange for the addition of a cryopump), the locations of the slits (dimensions given in the section entitled "Vacuum Chamber Components"), and approximate dimensions. The dashed lines show where the device may be sealed off to exclude unused volume. Again, as with the Wien filter, the device will likely require shielding with mu metal (Ref. 41, 42). The radius of curvature of ions entering the device is given by

$$R = (144/B\beta) (mE)^{1/2}$$

where  $E$  is the energy of the entering ions,  $B$  is the magnetic field,  $m$  is the ion mass, and  $\beta$  is the charge state (+, -). The resolution<sup>40</sup> is given by

$$(\Delta m/m) = w/[R [1-\cos(a)] + x \sin(a)]$$

where,  $w$  is the slit width,  $R$  is the radius of curvature, and  $x$  and  $a$  are defined in Fig. 16. Once again, a resolution of 0.01 is considered adequate.




Figure 16. Diagram of the Sector Magnet Showing Configuration and Construction Design.

Since the field produced by the superconducting magnet will be quite strong, a calculation was performed to yield an expression for its variation with distance on axis (Ref. 42). It is necessary to determine how far the source chain, and other devices (such as detectors) must be located to be safe from significant perturbations. The expression for the axial magnetic field is

$$B_z = [(B_0 (a^2 + \frac{L^2}{4})^{1/2}) / L] \{ (z + L/2) / (a^2 + (z + L/2)^2)^{1/2} - (z - L/2) / (a^2 + (z - L/2)^2)^{1/2} \},$$

where  $B_0$  is the field at the center of the magnet,  $a$  is the radius of the magnetic coils,  $L$  is the length of the magnet, and  $z$  is the axial distance from the center of the magnet. Plots of  $B_z/B_0$  are shown (linear, followed exponential in  $B$ ) in Figs. 17 and 18. Notice that for  $a=20$  cm and  $L=20$  cm,




Figure 17. Linear Plot of Magnetic Field Variation Along the Axis of the Superconducting Magnet (Normalized to the Central Field.)




Figure 18. Log Plot of Magnetic Field Variation Along the Axis of the Superconducting Magnet (Normalized to the Central Field).

the axial magnetic field is down by only three orders of magnitude at  $z=2$  meters (still  $\sim .08$  tesla).

#### Cluster Ion Detection

The detection system chosen depends upon the experiment. Since the initial experiments will measure radial density profiles and total charge, a Faraday cup with charge integrator will be used. The methodology for these measurements is well established in experiments done with pure electron plasmas (at UCSD) and with Lithium ion plasma experiments (at UCLA). The detector will be simply mounted on a standard 4 1/2 in. conflat flange and bolted to the top of the ion trap (Fig. 19).




Figure 19. Schematic of the Detection System Intended for Use in the First Experiments.

In this case the output of a radially movable Faraday cup is sent to a charge integrator. Calibration of the detector can be conveniently achieved by running the entire system in "free pass" mode, while using a known source flux. Another useful early experiment is to measure the total stored charge versus time as a function of cluster number. The Faraday cup will also suffice for these measurements.

Ultimately, wishing to determine the thresholds for the  $A^+$  and  $M^+$  growth processes (Refs. 14, 29), one must be able to discriminate the masses of the cluster ions. The steps of a typical experimental scenario would be these:

1. Load the trap with a known mass and number of cluster ions;
2. Add a known background density of reactant,  $H_2$  with known temperature for some time  $t$ ;
3. Analyze the products for total mass, and charge.

A direct approach to detection uses a setup like the one shown in Fig. 19. At the end of a drift tube is mounted an optional optics stage (not shown), followed by a sector magnet and channel plate detector. The product ions, ejected from the trap with well defined energy, are mass analyzed and detected with a channel plate in combination with a multichannel analyzer. This will give a signal that measures the relative numbers of clusters as a function of cluster mass. Hence, one can observe changes in population of different mass clusters in consecutive single measurements. The total charge can be determined by running the channel plate in total charge mode.

Alternatively, a more elegant method of detection (suggested by A. Y. Wong) uses a narrow band radio frequency source to excite selectively the cyclotron frequency of the ions, driving them to larger Larmor radii until they collide with and are collected on the trap inner surface. One simply attaches a charge integrator to the trap central wall and measures charge as a function of transmitter frequency.



Finally, well known nondestructive detection techniques have been in use for some time in pure electron plasmas (Ref. 43). In one method, the total number of charges can be determined by applying a potential to the end plates. The other method studies ion plasma waves and derives the plasma density.

#### The Cluster Ion Source

The importance of the cluster ion source to the success of this project should be stressed.

As mentioned, it is not clear that a cluster ion source exists that can fulfill all the requirements. The source should do the following:

1. Provide (a) very large cluster ions, (b) very low cluster temperature, and (c) very high flux;
2. Not use electron impact for ionization;
3. Not load the system with background gas;
4. Operate well around potentially large, stray magnetic fields.

The reasons behind requirements 1 (a) and (b) have been discussed previously. The stringency of requirement 1(b) can be estimated. The current,  $I$ , needed to fill the trap is given by

$$I = e J A = e n v A,$$

where  $J$  is the flux,  $e$  is the electron charge,  $n$  is the density of cluster ions in the trap,  $v$  is the velocity of the injected cluster ion beam, and  $A$  is the cross sectional area of the cluster ion plasma in the trap. For typical values (in cgs), one obtains

$$I = 10^{-19} \cdot 10^7 \cdot 10^8 \cdot 10^1 = 10^{-3} \text{ amps.}$$

If the total number of clusters arriving at the trap is only 1/10000 of the clusters generated by the source chain (recall  $\Delta m/m = 0.01$ ), then the instantaneous current needed at the source is 10 amps. Calculations indicate that the source need only be on for about 10 microseconds so the minimum estimated total output of the source is roughly  $10^{14}$  molecules of hydrogen.

Three concepts have emerged that could provide an adequate cluster ion source. To our knowledge, none of these sources, as described using hydrogen, have been reported in the literature. In Fig. 20(a), a standard pulsed gas valve is used to generate cluster ions that are simultaneously ionized by a suitable laser pulse. The only thing new here is the use of a laser to ionize the clusters. Since pulsed gas valves cannot be used at temperatures below which viton O-rings become brittle, high pressures of hydrogen gas must be used to achieve adequate cooling of the expanding gas for large cluster production. This necessitates differential pumping and skimming. The retardation grids are added to modify the residence time of the cluster ions. In Fig. 20(b), a solid surface of hydrogen is laser evaporated and laser ionized to produce an intense burst of clusters and ions. This method has been reported to generate cluster ions of carbon, and it has the advantages of low gas load and simplicity. In Fig. 20(c) a refrigerator is used to maintain a bath of liquid hydrogen. A pulse of hydrogen is introduced directly into the liquid from a pulsed valve that is thermally insulated from the cold bath. Finally, the expansion is ionized by a laser pulse. This source should produce high fluxes of very large cluster ions. The use of an inadequate cluster ion source will not prevent the experimental apparatus from working, but can limit its usefulness in studying CC and thus deserves careful attention.

#### System Performance

This section presents the results of calculations of maximum expected system performance (and plasma parameters) as a function of the cluster number (number of protons in the cluster ion) for the trap. The results presented here are intended to help clarify system capabilities in terms of




Figure 20. Proposed Cluster Ion Sources. (a) Pulsed gas valve/laser cluster ion source; (b) Solid surface/laser cluster ion source; (c) Pulsed gas valve evaporation/laser cluster ion source.

scalability to large sizes. The trap parameters used here are nearly the same as those of the proposed device, with the exception of length (the proposed device is 20 cm long). The parameters used in these calculations are

- B: The magnetic field (10 T).
- $\Phi$ : The potential applied to the central conductor (1000 Kvolts).
- a: The radius of the central conductor (0.3 cm).
- d: The radius of the trap wall (5 cm).
- r: The radius of the plasma (2.5 cm).
- $\Delta$ : The thickness of the (annular) plasma (0.2 cm).
- L: The length of the magnet (100 cm).

Fig. 21 shows the variation of the cyclotron frequency with cluster number. The cyclotron frequency is given by

B = 100kG,  $\Phi$  = 1000k-volts, a = 0.3cm, d = 5cm, r = 2.5cm,  $\Delta$  = 0.2cm, L = 100cm




Figure 21. Cyclotron Frequency Versus Cluster Number.

$$W_c = eB/mc,$$

where  $e$  is the charge,  $B$  is the magnetic field,  $m$  is the cluster mass, and  $c$  is the speed of light. Notice that it is linear and is only about 100 Hz for clusters of size number  $10^7$ .

Fig. 22 is obtained from the density limit relation (Ref. 44),

$$W_c^2 - (4 e \Phi) / (m R^{+2} \ln(d/r)) = W_p^2 [ \{ (8 R \Delta) / R^{+2} \} - \{ (4 R \Delta - 2 R^{+2} \ln(R^+/r) + 2 R^{-2} \ln(R^-/r)) / (R^{+2} \ln(d/r)) \} ].$$

In Fig. 22 the quantity

$$\Phi = \frac{4 e \phi}{M R^{+2} \ln(d/r)}$$

$B = 100 \text{ kG}$ ,  $\Phi = 1000 \text{ k-volts}$ ,  $a = 0.3 \text{ cm}$ ,  $d = 5 \text{ cm}$ ,  $r = 2.5 \text{ cm}$ ,  $\Delta = 0.2 \text{ cm}$ ,  $L = 100 \text{ cm}$




Figure 22. Central Rod Potential Versus Cluster Number at Constant Total Mass.

is plotted versus cluster number. This quantity is indicative of confinement potential of the bias (1000 KV) on the central conduction and has units of  $\text{sec.}^{-2}$ .

Definitions of the variables are the following:  $\omega_c$  is the cyclotron frequency,  $e$  is the charge,  $\Phi$  is the potential on the central rod,  $m$  is the mass,  $R^+$  is the outside radius of the plasma,  $d$  is the radius of the trap,  $r$  is the radius of the plasma (to its center), twice  $\Delta$  is the width of the plasma, and  $R^-$  is the inside radius of the plasma. The linear decrease is just the result of less space charge repulsion per mass as the size of the clusters increases.

Fig. 23 shows the variation in the characteristic plasma frequency with increasing cluster number (at constant trapped mass). The plasma

$B = 100\text{kG}$ ,  $\Phi = 1000\text{k-volts}$ ,  $a = 0.3\text{cm}$ ,  $d = 5\text{cm}$ ,  $r = 2.5\text{cm}$ ,  $\Delta = 0.2\text{cm}$ ,  $L = 100\text{cm}$




Figure 23. Plasma Frequency Versus Cluster Number.

frequency is given by

$$\omega_p^2 = (4 \Pi n e^2) / (\epsilon_0 m),$$

where,  $\Pi$  is 3.14,  $n$  is the plasma density,  $e$  is the cluster charge,  $\epsilon_0$  is the permittivity of free space, and  $m$  is the cluster mass. Notice that at a cluster size near  $10^6$  the plasma frequency reaches a minimum plateau. The plateau indicates the increasing dominance of electrostatic confinement for high cluster number. It also indicates a balance between the number of charges on the central conductor and the number of charges in the plasma.

Fig. 24 is a plot of the confinable number density that may be contained in the trap for the given magnetic field and central conductor bias. Note that the density needed begins to remain constant at about  $10^6 \text{ cm}^{-3}$  for mass numbers above  $10^7$ . Fig. 25 is the same as 24, except given in terms of mass per volume.

Fig. 26 is a plot of the total confinable mass in the trap as a function of the cluster number. Notice that, with large cluster sizes (of around  $10^6$ ), the total stored mass increases sharply.

Fig. 27 is the same as 26 except with a much larger trap. The trap parameters in this case are

- B: The magnetic field (100 T).
- $\Phi$ : The potential applied to the central conductor (1000 Kvolts).
- a: The radius of the central conductor (50.0 cm).
- d: The radius of the trap wall (100 cm).
- r: The radius of the plasma (80.0 cm).
- $\Delta$ : The thickness of the (annular) plasma (6.0 cm).
- L: The length of the magnet (200.0 cm).

Notice that with this (only somewhat futuristic) trap, the storage of greater than 0.3 mg of cluster ions is possible in a device only two meters long. This corresponds to an energy density of  $5 \times 10^6 \text{ J/m}^3$ .

B = 100kG, Phi = 1000k-volts, a = 0.3cm, d = 5cm, r = 2.5cm, del = 0.2cm, L = 100cm




Figure 24. Plasma Number Density Versus Cluster Number.

B = 100kG, Phi = 1000k-volts, a = 0.3cm, d = 5cm, r = 2.5cm, del = 0.2cm, L = 100cm




Figure 25. Mass Density Versus Cluster Number.



$B = 100\text{kG}$ ,  $\Phi = 1000\text{k-volts}$ ,  $a = 0.3\text{cm}$ ,  $d = 5\text{cm}$ ,  $r = 2.5\text{cm}$ ,  $\text{del} = 0.2\text{cm}$ ,  $L = 100\text{cm}$




Figure 26. Total Mass Confined by the Trap Versus Cluster Number for a 100-cm Trap. (Note the dramatic increase beyond cluster numbers of  $10^7$ .) This trap is 5 times longer than the one proposed; the remaining parameters are similar.

$B = 1000\text{kG}$ ,  $\Phi = 1000\text{k-volts}$ ,  $a = 50\text{cm}$ ,  $d = 100\text{cm}$ ,  $r = 80\text{cm}$ ,  $\text{del} = 6\text{cm}$ ,  $L = 200\text{cm}$




Figure 27. Total Mass Confined by the Trap Versus Cluster Number for a 300-cm "Next-Generation" Trap. (Note the dramatic increase beyond cluster numbers of  $10^7$ .) This trap is 10 times longer and the field is higher than the one proposed; the remaining parameters are similar.

## CONCLUSIONS AND RECOMMENDATIONS

In conclusion, the research effort for the past two years has resulted in the development of the CI approach to CC (Containerless Condensation). The general strategy in the years ahead is to develop CC techniques, more specifically, to build the first pseudo-wall designed for long-term studies of very large hydrogen cluster ions. If successful, this will fulfill the near-term goal of this project: Implementation of the experimental investigation and understanding the formation of hydrogen cluster ions from the viewpoint of "containerless" condensation.

Recommendations pertaining to the construction and operation of the cluster ion trap have been made in the previous section. It is anticipated that the program will require one year to construct and test the basic apparatus. The first experiments (previously stated) should start the second year. The remainder of this section gives an approximate timetable covering the next three years.

The effort for the first year will involve the purchase, installation, and checkout of the UHV system. This should include all of the components given in Fig. 10. The Wien filter and sector magnet should also be built by the appropriate subcontractors. One of the two optional cryopumps could be purchased and attached either at the 6-way cross or the sector magnet.

Professor A. Y. Wong and Dr. G. Rosenthal should be consulted on matters concerning system construction. Professor Wong and Dr. Rosenthal have agreed to set aside part of their time this year for this purpose.

Also, during the first year, needed experimentation on the cluster ion source should be carried out. Low background flux, pulsed sources should be employed exclusively. Initially, a commercial source manufactured by Coultron was under consideration, but this was abandoned because of the need for too many modifications and limited mass/charge handling capability.

Some construction on the ion trap should also begin the first year.

The cryopump is the final stage of pumping, used for obtaining very high vacuum at high pumping speed.

The second year should involve the construction and implementation of the cluster ion source and ion trap. The source and most of its components should be constructed and tested on site. The ion trap should be constructed by an appropriate vendor (suggest UCLA machine shop) under the care and supervision of persons with first-hand experience. The surfaces will be machined to high tolerance from a single supply of oxygen-free copper. Special care is required in the mounting and assembly to avoid field inhomogeneities, junction potentials, and stray capacitances. The trap inner sections could also be electroformed. This is more expensive, but yields a better surface.

The superconducting magnet (8.5 Tesla) should be custom made by an experienced commercial vendor (recommend Oxford). They can also provide the liquid helium baffle and all of the supporting equipment for the magnet.


The detection hardware should be purchased and assembled on site during the second year. If all goes well, the system should be ready for checkout near the end of the second year.

With the system near completion, the third year will involve performing the first experiments. As mentioned previously, these will measure the plasma radial density profiles, the total charge in the trap, and the confinement times as a function of cluster size.


In later experiments, the ion trap will be filled with cluster ions of known size and temperature. Reactant (cold  $H_2$  or H) will be added to the cooled cluster ion pseudo-wall. Cooling cycles may be needed during the addition of the reactants. After some time, the ion cloud composition will be analyzed by releasing the trap contents for mass and energy analysis. The cycle is then repeated. Cluster ion growth rates and critical information are then determined from the data.

In the fragmentation experiments, the scenario is similar. The ion trap is again loaded with cluster ions of known composition and temperature. They are then exposed to an appropriate collider (protons, antiprotons, positrons, photons, etc.) of well defined energy. The composition that remains in the trap is then mass and energy analyzed. The results of the fragmentation studies will decide if large cluster ions can be conveniently fragmented into just a few large charged pieces (with minimum H or H<sub>2</sub> loss).

A series of experiments should be done that involve the multiplication of seed crystals by fragmentation. The general fragmentation reaction using protons is



The general fragmentation reaction for photons is



where k and n are integers. Note that the resultant fragments must remain captured and cooled. The first fragmentation reaction has the highest priority.

## REFERENCES

1. Forward, R. L., Antiproton Annihilation Propulsion, AFRPL TR-85-034, Forward Unlimited/University of Dayton Research Institute, Dayton, Ohio, September 1985.
2. Forward, R. L., Prospects for Antihydrogen Production, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 9-26, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
3. Be, S. H., Yano, K., Enjoji, H., and Okamoto, K, "Necessary Conditions for Generating Intense Cluster Beam," Jap. J. of Appl. Phys., Vol. 14, No. 7, pp. 1049-1056, January 1975.
4. Yano K., and Be, S. H., "Mass Analysis of Cluster Ion Beams by Wien Filter," Jap. J. of Appl. Phys., Vol. 19, No. 6, pp. 1019-1025, January 1980.
5. Wineland, D. J., Ion Traps for Large Storage Capacity, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 181-194, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
6. Barlow, S. E., The Dynamics of Stored Ions and Low Temperature Ion-Molecule Reactions, Ph.D. Dissertation, Physics Department, University of Colorado, Boulder, CO.
7. Carrington, A., and Kennedy, R. A., "Infrared Predissociation of the  $H_3^+$  Ion," J. Chem. Phys., Vol. 81, No. 1, pp. 91-112, July 1984.
8. Crofton, M., Infrared Spectroscopy of  $H_3^+$ , Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8- 1/9/87, pp. 121-124, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.

9. Yeh, L., Hydrogen Cluster Ion Spectroscopy Obtained through Vibrational Predissociation, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 127-141, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
10. Bowers, M., Formation and Reactivity of Small Hydrogen Cluster Ions, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 63-72, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
11. Schaefer, H. III,  $H_n^+$  Radiative Properties, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 75-79, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
12. Saxon, R., Overview of Hydrogen Clusters, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 29-37, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
13. Michels, H., and Montgomery, J., Electronic Structure and Stability of Small Cation and Anion Hydrogen Cluster Ions, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 83-94, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
14. Stwalley, W. C., Large Hydrogen Cluster Ions, Proc. Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8- 1/9/87, pp. 41-45, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
15. Bahns, J. T., Sando, K. M., Tardy, D. C., and Stwalley, W. C., Proc. of the Hydrogen Cluster Ion Workshop, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop," 1/8-1/9/87, Appendix pp. 1-70, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
16. Saxon, R. P., and Talbi D., Excited States of Hydrogen Cluster Ions, University of Dayton Research Institute, Subcontract Order RI-49857X, October 1987.

17. Helm, H., Photon-Assisted Formation and Cooling of Molecular Hydrogen, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 97-108, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
18. Bae, Y. K., Coggiola, M. J., and Peterson, J. R., "Search for  $H_2^-$ ,  $H_3^-$ , and other Metastable Negative Ions," Phys. Rev. A, Vol. 29, p., 2888, 1984.
19. Neumann, R., Poth, H., Winnacker, A., and Wolf, A., "Laser Enhanced Electron-Ion Capture and Antihydrogen Formation," Z. Phys. A-Atoms and Nuclei, Vol. 313, p. 253, 1983.
20. Gabrielse, G., First Capture of Antiprotons in an Ion Trap and the Possibility of Antihydrogen, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 167-177, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
21. McIlrath, T. J., Long Duration Coherent Lyman-Alpha Sources, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 197-203, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
22. Phillips, W., Gould P., and Lett P., Electromagnetic Manipulation of Atomic Hydrogen, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 159-164, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
23. Lovelace, R., Magnetic Traps for an Anti-Hydrogen Gas, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 207-217, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
24. Cline, R. W., Smith, D. A., Greytak, T. J., and Kleppner, D., "Magnetic Confinement of Spin-Polarized Atomic Hydrogen," Phys. Rev. Lett., Vol. 45, No. 26, pp. 2117-2120, December 1980.

25. Stwalley, W. C., "A Hybrid Laser-Magnet Trap for Spin-Polarized Atoms," Prog. Quant. Electr., Vol 8, pp. 203-208, 1984.
26. Weiner, J., Associative Two-Body Condensation in Laser-Cooled Sodium and Hydrogen Cluster Ions, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 49-59, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
27. Lester, W., and Huang, S., Quantum Monte Carlo Study of the MIES Associated with  $H_2(X^1\Sigma_g^+)$  and  $H_2(B^1\Sigma_u^+)$ , Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 111-118, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
28. Stwalley, W. C., Bibliography of Hydrogen Cluster Ions, Paper presented at the Second Antiproton Science and Technology Workshop, RAND Corporation, Santa Monica, CA, October 1987.
29. Stwalley, W. C., The Synthesis of Large Cluster Ions from Elementary Constituents: A Possible Route to Bulk Antimatter, Paper presented at the Second Antiproton Science and Technology Workshop, RAND Corporation, Santa Monica, CA, October 1987.
30. Casero, R. and Soler, J. M., personal communication.
31. Echt, O., Multiply Charged Clusters, The Physics and Chemistry of Small Clusters, NATO ASI Series, Ed. P. Jena, B. K. Rao, and S. N. Khanna, 1986.
32. Driscoll, C. F., Containment of Single-Species Plasmas at Low Energies, Low Energy Antimatter, Ed. D. B. Cline, World Scientific, Singapore, 1986.
33. Malmberg, J. H., O'Neil, T. M., Hyatt, A. W., and Driscoll, C. F., The Cryogenic Pure Electron Plasma, Proc. 1984 Sendai Symposium on Plasma Nonlinear Phenomena.



34. Brown, L. S., and Gabrielse G., "Geonium Theory: Physics of a Single Electron or Ion in a Panning Trap," Rev. Mod. Phys., Vol. 58, No. 1, pp. 233-311, January 1986.
35. Mitchell, J., The Role of Electron-Ion Recombination in Bulk Antimatter Production, Proc. Cooling Condensation and Storage of Hydrogen Cluster Ions Workshop, 1/8-1/9/87, pp. 1435-156, Ed. J. T. Bahns, University of Dayton Research Institute, 1987.
36. Stwalley, W. C., Private communication, obtained from Relativistic Particle Properties Data Base, 1984. Figures courtesy of V. Flaminio, W. G. Moorhead, D. R. O. Morrison, and N. Rivoire, CERN.
37. Kluge, H. J., Schnatz, H., and Schweikhard, L., "Penning Trap for Studying Cluster Ions," Phys. D--Atoms Molecules and Clusters, Vol. 3, pp. 189-194, 1986.
38. Rosenthal, G., Dimonte, G., and Wong, A. Y., "Stabilization of the Diochotron Instability in an Annular Plasma," Phys. Fluids, Vol. 30, No. 10, pp. 3257-3261, October 1987.
39. O'Hanlon, J. F., A User's Guide to Vacuum Technology, John Wiley and Sons, New York, NY, 1980.
40. Moore, J. H., Davis, C. C., and Coplan, M. A., Building Scientific Apparatus, Addison-Wesley Publishing Co., Cleveland, OH, 1983.
41. CRC Handbook of Chemistry and Physics, 55th Ed., CRC Press, Cleveland, OH, 1975, p. E-120.
42. Jackson, J. D., Classical Electrodynamics, John Wiley & Sons, New York, NY, 1975.
43. deGrassie, J. S., and Malmberg, J. H., "Wave-Induced Transport in the Pure Electron Plasma," Phys. Rev. Lett., Vol. 39, No. 17, pp 1077-1080, October 1977.

44. Rosenthal, G., Wong, A. Y., and Simon, M., "Confinement of High Density Pure Ion Plasmas," to be submitted to Phys. Rev. Lett., 1988.
45. Davidson, R. C., Theory of Nonneutral Plasmas, Benjamin, Publishing Co., Reading MA, 1974.

## APPENDIX A

### Radial Confinement and the Classical Brillouin Limit

The presented derivation may be found in R. C. Davidson's book (Ref. 45).

For the confinement, a charged particle undergoing uniform circular motion in a stable orbit that lies in a region within electric and magnetic fields, the centrifugal force must balance the Lorentz force (see Fig. A-1),

$$-\frac{m v^2}{r} + e E_r^0 + v_\theta^0 B = 0 \quad (\text{A.1})$$

where  $E_r^0$  and  $v_\theta^0$  are the radial electric field and azimuthal velocity at equilibrium,  $m$  is the mass,  $r$  is the orbit radius, and  $B$  is the magnetic field.

Since the radial electric field is self-generated by the plasma, it must be determined self-consistently using Poisson's equation,

$$\frac{1}{r} \frac{d}{dr} (r E_r^0) = -4 \pi e n^0(r) \quad (\text{A.2})$$

The simplest solution is to take  $n^0(r)$  to be constant; then the electric field may be obtained through direct integration giving

$$E_r^0 = -r 4 \pi e \frac{\bar{n}}{2} \quad , \quad (\text{A.3})$$

for  $0 < r < R_p$ . This may be expressed in terms of the plasma frequency

$$\omega_p^2 = \frac{4 \pi \bar{n} e^2}{m} \quad , \quad (\text{A.4})$$




Figure A.1. Plot of the Two Rotational Frequencies of a "Solid" Plasma and the Location of the Brillouin Limit. Below are shown the conventions used in the derivation presented in the text (Appendix).

as

$$E_r^0 = \frac{m \omega_p^2 r}{2 e} \quad (\text{A.5})$$

for  $0 < r < R_p$ . Introducing the angular frequency

$$\omega_v = \frac{v_\theta^0}{r}, \quad (\text{A.6})$$

and the cyclotron frequency,

$$\Omega = \frac{e B_0}{m c}, \quad (\text{A.7})$$

into equation (A.1), it may be expressed as

$$\omega^2 + \Omega \omega + \frac{\omega_p^2}{2} = 0. \quad (\text{A.8})$$

The solutions to this quadratic are

$$\omega = \omega^\pm = \frac{\Omega}{2} \left[ 1 \pm \sqrt{1 - \frac{2 \omega_p^2}{\Omega^2}} \right]^{\frac{1}{2}}. \quad (\text{A.9})$$

The two rotational frequencies  $\omega^+$  and  $\omega^-$  are plotted in the figure as a function of  $\frac{2 \omega_p^2}{\Omega^2}$ .

In the low density limit the fast rotation,  $\omega^+$ , corresponds to all the particles rotating about the axis of symmetry at the cyclotron frequency. The low frequency,  $\omega^-$ , corresponds to the trochoidal orbits which give rise to a slow  $E^0 \times B_0 e_z$  rotation of the plasma.

The high density limit is known as the Brillouin limit. It can be seen from Equation (A.9) that when  $\frac{2 \omega_p^2}{\Omega^2} = 1$ , then

$$\omega^+ = \omega^- = \frac{\Omega}{2}, \quad (\text{A.10})$$

Furthermore, for the solutions to Equation (A.9) to be real, we must have,

$$\omega_p^2 < \frac{\Omega^2}{2} . \quad (A.11)$$

This equation states that the electrostatic repulsive forces, measured by  $\omega_p^2$  must be overcome by the magnetic restoring force, measured by  $\Omega^2$ . Inserting the definitions for  $\omega_p^2$  and  $\Omega^2$  into Equation (A.11) yields the relation

$$n m c^2 < \frac{B^2}{8\pi} , \quad (A.12)$$

where  $n$  is the density,  $m$  is the mass of the cluster ion, and  $B$  is the magnetic field. This equation gives the limit to the mass density ( $nm$ ) that can be achieved under the assumption of a solid cylindrical cylinder with constant radial density. This limit may be exceeded in two ways:

1. By operating in the low density ( $n$ ), high mass ( $m$ ) regime (see plots in the section entitled "System Performance").
2. By using an annular plasma, neutralized by a central conducting rod.<sup>37,43</sup>

#### System Pumping

As discussed in the section of the report entitled "The Pumping System" (see Fig. 14a) the pressure in the trap is given by

$$P_t = \frac{P_s}{1 + \frac{c_2}{c_1} \frac{s_p}{s_p + c_2}} , \quad (A.13)$$

where  $P$  is the pressure in the source or trap (Pa or Torr),  $C$ 's are the conductances (L/s), and  $S$  the speed of the pump (L/s). We want to include the outgassing rate of the baked stainless steel ( $10^{-13}$  W/m<sup>2</sup>) in the derived expression for the pumping speed.

Define

$$s_p' = \frac{s_p c_2}{s_p + c_2} , \quad (A.14)$$

then

$$P_t = \frac{P_s}{1 + \frac{s_p}{c_1}} , \quad (A.15)$$

or

$$P_t = \frac{c_1 P_s}{c_1 + s_p} . \quad (A.16)$$

Now,  $P=Q/C$ , and applying this to the source and trap gives

$$P_s - P_t = \frac{Q_1 + Q_w}{c_1} , \quad (A.17)$$

rearranging gives,

$$c_1 P_s = c_1 P_t + Q_1 + Q_w ; \quad (A.18)$$

inserting this into equation (A.16) gives

$$P_t = \frac{c_1 P_t + Q_1 + Q_w}{c_1 + s_p} ; \quad (A.19)$$

solving this expression for  $P_t$  gives

$$P_t = \frac{Q_1 + Q_w}{s_p} , \quad (A.20)$$

$$= \frac{c_1 (P_s - P_t) + \alpha_w A_w}{s_p} \quad (A.21)$$

$$= \frac{\alpha_w A_w + c_1 P_s}{s_p + c_1} \quad (A.22)$$

$$= \frac{\alpha_w A_w + c_1 P_s}{c_1 + \frac{s_p c_2}{c_2 + s_p}} \quad (\text{A.23})$$

$$= \frac{(c_2 + s_p) (\alpha_w A_w + c_1 P_s)}{c_1 (c_2 + s_p) + s_p c_2} \quad (\text{A.24})$$

$$= \frac{(c_2 + s_p) (\alpha_w A_w + c_2 P_s)}{c_1 c_2 + s_p (c_1 + c_2)} \quad (\text{A.25})$$

Now, solving this expression for  $s_p$  gives

$$s_p = \frac{(c_2 + s_p) (\alpha_w A_w + c_1 P_s) - c_1 c_2 P_t}{P_t (c_1 + c_2)} \quad (\text{A.26})$$

The strategy in the system design is to make the conductance of the source chain ( $c_1$ ) very small relative to  $c_2$  (which is made very big), and this can be achieved to a good approximation in the design presented. Thus, taking the appropriate limit gives

$$s_p \lim_{c_2} = \frac{\alpha_w A_w + c_1 P_s - c_1 P_t}{P_t} \quad (\text{A.27})$$

Specifying the parameters on the right side as

$$\alpha_w = 10^{-13} \frac{W}{M^2} \quad (\text{A.28})$$

$$A_w = 1.26 M^2 \quad (\text{A.29})$$

$$P_s = 10^{-9} \text{ Torr} \quad (\text{A.30})$$

$$P_t = 10^{-11} \text{ Torr} \quad (\text{A.31})$$

$$c_1 = 121 \frac{d^3}{L} = .054 \frac{M^3}{s} \quad (\text{A.32})$$

This finally gives the pumping speed as



$$S_p = \frac{1.26 \times 10^{-10} + 7.18 \times 10^{-6}}{1.33 \times 10^{-9}} = 54. \frac{L}{s} \quad (A.33)$$

$$S_p = 5344 \cdot \frac{L}{s} \quad (A.34)$$

#### Transit Spreading of the Beam

This "worst case scenario" calculation assumes that a cluster ion plasma of density  $n_o$ , mass  $m$ , charge  $q$  is formed instantaneously and then expands in the  $x$  direction under the influence of the self electric field,  $E$ . The equation of motion is

$$\begin{aligned} m x &= q E \\ x &= \frac{q E}{m} \end{aligned} \quad (A.35)$$

Inserting the electric field,  $E = (mR/2q) \omega_p^2$ , gives

$$\begin{aligned} a = x &= \frac{q m R 4 \pi q^2 n_o}{m 2 q m} = \frac{2 \pi q^2 n_o R}{m} \\ a &= \frac{\omega_p^2 R}{2} \end{aligned} \quad (A.36)$$

where the plasma frequency is given by

$$\omega_p^2 = \frac{4 \pi q^2 n_o}{m} \quad (A.37)$$

The Newtonian equations of motion are

$$\begin{aligned} v &= v_o + a t \\ v &= x_o + v_o t + \frac{1}{2} a t^2 \end{aligned} \quad (A.38)$$

The initial conditions are

$$v_o = 0 \text{ and } x_o = R_i \quad (A.39)$$

Where  $R_i$  and  $R_f$  are the radii of the source and the ion trap respectively,

$$R_f = R_i + \frac{1}{2} a t^2 \quad (\text{A.40})$$

$$(R_f - R_i) \frac{2}{t^2} = \frac{\omega_p^2 R_i}{2} \quad (\text{A.41})$$

solving for t gives

$$t^2 = \frac{4 (R_f - R_i)}{R_i \omega_p^2} \quad (\text{A.42})$$

$$t = 2 \frac{R_f - R_i}{R_i}^{\frac{1}{2}} \frac{1}{4 \pi}^{\frac{1}{2}} \frac{m}{q n_o}^{\frac{1}{2}} \quad (\text{A.43})$$

This equation gives the maximum allowable transit time for the beam of cluster ions. It may be recast into a more convenient form, by defining  $2\alpha$  as the mass of a cluster divided by the mass of a proton,  $z$  as number of proton charges per cluster, and  $n_i$  as the number density of cluster ions ( $\text{cm}^{-3}$ ).

$$t = 1.52 \times 10^{-3} \frac{R_f - R_i}{R_i}^{\frac{1}{2}} \frac{1}{z} \frac{2\alpha}{n_i} \quad (\text{A.44})$$

The constant in front has the dimensions ( $\text{s cm}^{3/2}$ ). Typical values for  $R_i$  and  $R_f$ , the cluster sizes, and densities give times in the milliseconds. The length the cluster ions will travel is roughly 2 meters, implying a cluster ion injection velocity of  $2 \times 10^5 \text{ cm s}^{-1}$ . This is easily achieved.

## APPENDIX B

### Bibliography of Hydrogen Cluster Ions

William C. Stwalley

Center for Laser Science and Engineering and  
Departments of Chemistry and of Physics and Astronomy,  
University of Iowa, Iowa City, Iowa 52242-1294

## Introduction

The enclosed bibliography includes the edited results of September 1987 Chemical Abstracts On-Line Searches, plus selected additional references, for the period 1966-1987 for the species  $H_2^+$  (and  $H_2^-$ ),  $H_3^+$  (and  $H_3^-$ ), and  $H_N^+$  and ( $H_N^-$ ,  $N \geq 4$ ), in Sections I, II, and III, respectively. References are roughly in reverse chronological order. Note that  $H_2^-$  is not a stable species and the evidence suggests  $H_3^-$  is also unstable. All references have been copied or requested through interlibrary loan. The assistance of Todd Colin and Tom Yang in these searches and partial support from the Rand Corporation are gratefully acknowledged.

Section I:  $H_2^+$  (and  $H_2^-$ )

1. Sen, Amarjit; McGowan, J. W.; Mitchell, J. B. A., "Production of Low-Vibrational-State Hydrogen Ions ( $H_2^+$ ) for Collision Studies", J. Phys. B: At. Mol. Phys. 20(7), 1509-15 (1987).
2. Wolniewicz, L.; Poll, J. D., "On the Higher Vibration-Rotational Levels of Hydrogen Deuteride Ion ( $HD^+$ ) and Hydrogen Molecular Ion ( $H_2^+$ )", Mol. Phys. 59(5), 953-64 (1986).
3. Stine, J. R.; Muckerman, J. T., "Critical Evaluation of Classical Trajectory Surface-Hopping Methods as Applied to the Hydrogen Molecular Ion ( $H_2^+$ ) + Hydrogen Molecule ( $H_2$ ) System", J. Phys. Chem. 91(2), 459-66 (1987).
4. Helm, H.; Cosby, P. C.; "Photodissociation Measurement of Rovibrational Energies and Populations of Molecules in Fast Beams", J. Chem. Phys. 86(12), 6813-22 (1987).
5. Xu, E. Y.; Helm, H.; Kachru, R., "Field Ionization of High-Lying States of  $H_2$ ", Phys. Rev. Lett. 59(10), 1096-99 (1987).
6. Niedner, G.; Noll, M.; Toennies, J. P.; Schlier, Ch., "Observation of Vibrationally Resolved Charge Transfer in  $H^+ + H_2$  at  $E_{CM} = 20$  eV", J. Chem. Phys. 87(5), 2685-2694 (1987).
7. Schlier, Christoph G.; Vix, Ulrike, "Complex Formation in Proton-Hydrogen Collisions. II. Isotope Effects", Chem. Phys. 113, 211-221 (1987).
8. Weiner, J., "Associative Two-Body Condensation in Laser-Cooled Sodium and Hydrogen", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 47-59. Edited by John T. Bahns. University of Dayton Research Institute (1987).
9. Helm, H., "Photon-Assisted Formation and Cooling of Molecular Hydrogen", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 95-108. Edited by John T. Bahns. University of Dayton Research Institute (1987).
10. Mitchell, J. B. A., "The Role of Electron-Ion Recombination in Bulk Antimatter Production", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 143-156. Edited by John T. Bahns. University of Dayton Research Institute (1987).
11. Bahns, J. T., "Key Problems and Hydrogen Atom Formation", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 219-230. Edited by John T. Bahns. University of Dayton Research Institute (1987).

12. Bahns, J. T.; Sando, K. M.; Tardy, D. C.; Stwalley, W. C., "Proceedings of the Hydrogen Cluster Ion Study Group", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, A1-64. Edited by John T. Bahns. University of Dayton Research Institute (1987).
13. De Bruijn, D. P.; Neuteboom, J.; Govers, T. R.; Los, J., "Dissociative Decay of  $N = 3$  Levels in  $H_3$ . I. Populated in Charge Exchange of Hydrogen Ion ( $H_2^+$ ) with Cesium", Phys. Rev. A: Gen. Phys. 34(5), 3847-54 (1986).
14. Richards, J. A.; Larkins, F. P., "Photoionization Cross Section Calculations of Molecular Hydrogen and Hydrogen Ion ( $H_2^+$ ) Using Numerical Continuum Wavefunctions", J. Phys. B: At. Mol. Phys. 19(13), 1945-57 (1986).
15. Chandra, N., "Ab Initio Multichannel Photoionization of Molecular Hydrogen: Photoelectron Angular Distribution for Rotationally Resolved States of Hydrogen Ion ( $H_2^+$ )", J. Phys. B: At. Mol. Phys. 19(13), 1959-88 (1986).
16. Blaise, Paul; Henri-Rousseau, Olivier, "On the Quantum Correlation Between the Kinetic and Potential Energies in the Hydrogen Molecular Ion ( $H_2^+$ )", C. R. Acad. Sci. Ser. 2, 302(6), 297-302 (1986).
17. Stehle, C.; Feautrier, N., "Absorption or Emission During a Collision: A Test Case Hydrogen Molecular Ion ( $H_2^+$ )", J. Phys. (Les Ulis, Fr.), 47(6), 1015-20 (1986).
18. Miao, Jingwei; Yang Beifang; Hao, Shizuho; Jiang, Zengxue; Shi, Miangong; Cue, N., "Internuclear Separations From Foil Breakup of Fast  $H_2^+$ ,  $H_3^+$ ,  $D_2^+$  and  $D_3^+$  Molecules", Nucl. Instrum. Methods Phys. Res. Sect. B, B13(1-3), 181-3 (1986).
19. Blaise, Paul; Henri-Rousseau, Olivier, "Accuracy of Approximate Wave Functions for the Molecular Hydrogen Ion ( $H_2^+$ ): Calculation of Standard Deviations for the Hamiltonian in the  $2\Sigma_g^+$  State", C. R. Acad. Sci. Ser. 2, 301(13), 907-10 (1985).
20. Cizek, Jiri; Damburg, R.; Graffi, Sandro; Grecchi, Vincenzo; Harrell, Evans M., II; Harris, Jonathan G.; Nakai, Sachiko; Paldus, Josef; Propin, R.; Silverstone, Harris J., "1/R Expansion for Molecular Hydrogen Ion ( $H_2^+$ ): Calculation of Exponentially Small Terms and Asymptotics", Phys. Rev. A: Gen. Phys. 33(1), 12-54 (1986).
21. Liao, C. L.; Ng, C. Y., "Vibrational State Distributions of  $H_2^+$  ( $v$ ) Resulting From the Hydrogen Ion-Molecule Electron Transfer Reactions  $H_2^+(v_0' = 0, 1) + H_2(v_0'' = 0) \rightarrow H_2(v') + H_2^+(v)$  in the Collisional Energy Range of 2-16 eV", J. Chem. Phys. 84(1), 197-200 (1986).
22. Sen, A.; Mitchell, J. B. A., "Production of Vibrationally Cold Ions Using

- a Radio-Frequency Storage Ion", Rev. Sci. Instrum. 57(5), 754-756 (1986).
23. Bonnie, J. H. M.; Eenshyistra, P. J.; Los, J.; Hopman, H. J., "Influence of the Vibrational Quantum Number of the Resonant State in Resonant Multiphoton Ionization/Dissociation of Hydrogen Molecules", Chem. Phys. Lett. 125(1), 27-32 (1986).
  24. Graffi, S.; Grecchi, V.; Harrell, E. M., II; Silverstone, H. J., "The 1/R Expansion for  $H_2^+$ : Analyticity, Summability, and Asymptotics", Ann. Phys. (N. Y.), 165(2), 441-83 (1985).
  25. Eaker, Charles W.; Muzyka, Jennifer L., "A Quasiclassical Trajectory Study of the Hydrogen-Deuterium  $((H_2 + D_2)^+)$  System", Chem. Phys. Lett. 119(2-3), 169-72 (1985).
  26. Primorac, M.; Kovacevic, K., "An Application to Molecular Hydrogen Ion  $(H_2^+)$  of Laplace Type Integral Transform and Its Inverse", Z. Naturforsch. A: Phys. Phys. Chem., Kosmophys. 40A(3), 246-50 (1985).
  27. Mohlmann, G. R., "Stimulated Raman Scattering from  $H_2^+$  Ions", Chem. Phys. Lett. 115(2), 226-229 (1985).
  28. Brenton, A. G.; Fournier, P. G.; Govers, B. L.; Richard, E. G.; Beynon, J. H., "The Vibrational Population Distribution of  $H_2^+$  Formed From a Series of Different Precursor Molecules", Proc. R. Soc. London, A, 395(1808), 111-25 (1984).
  29. Nicolaides, C. A.; Petsalakis, I. D.; Theodorakopoulos, G., "Theory of Chemical Reactions of Vibronically Excited Molecular Hydrogen ( $B^1\Sigma_u^+$ ). III. Formation of Bound Excited States of the Hydrogen  $(H_2)_2$ ,  $(H_2)_3$ , and  $(H_2)_5$  Clusters", J. Chem. Phys. 81(2), 748-53 (1984).
  30. Bae, Y. K.; Coggiola, M. J.; Peterson, J. R., "Search for  $H_2^-$ ,  $H_3^-$ , and Other Metastable Negative Ions", Phys. Rev. A, 29(5), 2888-90 (1984).
  31. Damburgs, R.; Propin, R. Kh.; Graffi, Sandro; Grecchi, Vincenzo; Harrell, Evans M., II; Cizek, Jiri; Paldus, Josef; Silverstone, Harris J., "1/R Expansion for Hydrogen Molecular Ion  $(H_2^+)$ : Analyticity, Summability, Asymptotics, and Calculation of Exponentially Small Terms", Phys. Rev. Lett. 52(13), 1112-15 (1984).
  32. Lee, Chyuan Yih; DePristo, Andrew E., "Semiclassical Investigation of Vibrational State and Molecular Orientation Effects in Electron Transfer Reactions for the Hydrogen  $(H_2^+/H_2)$  Collision", J. Chem. Phys. 80(3), 1116-26 (1984).
  33. Kutina, R. E.; Edwards, A. K.; Pandolf, R. S.; Berkowitz, J., "UV Laser Photodissociation of Molecular Ions", J. Chem. Phys. 80(9), 4112-9 (1984).

34. Lee, Chyuan Yih; DePristo, Andrew E., "A Simple Model for the Interaction Potentials in Electron-Transfer Reactions: Application to the Molecular Hydrogen Ion/Molecular Molecular ( $H_2^+/H_2$ ) System", J. Am. Chem. Soc. 105(23), 6775-81 (1983).
35. Semenova, N. V., "Quasiclassical Calculation of Low-Lying Terms of Hydrogen ( $H_2^+$ ) Ion", Vestn. Leningr. Univ., Fiz., Khim. 3, 72-6 (1983).
36. Samsonov, B. F., "Calculation of Diatomic Molecules by the MO LCAO Method. Excited States of Hydrogen Ion ( $H_2^+$ )", Izv. Vyssh. Uchebn. Zaved., Fiz. 26(8), 115-17 (1983).
37. Chu, Shih I; Laughlin, Cecil; Datta, Krishna K., "Two-Photon Dissociation of Vibrationally Excited Molecular Hydrogen Ion ( $H_2^+$ ). Complex Quasi-Vibrational Energy and Inhomogeneous Differential Equation Approaches", Chem. Phys. Lett. 98(5), 476-81 (1983).
38. Brenton, A. G.; Beynon, J. H.; Richard, E. G.; Fournier, P. G., "Rotational Predissociation of  $H_2^+$  Ions of Different Precursor Origins", J. Chem. Phys. 79(4), 1834-45 (1983).
39. Ozaki, Jiro; Tomishina, Yasuo, "Monte Carlo Solutions of Schroedinger's Equation for Molecular Hydrogen ( $H_2^+$ ) Ion in Strong Magnetic Fields. II", J. Phys. Soc. Jpn. 52(4), 1142-7 (1983).
40. Khersonskii, V. K., "Frequencies and Probabilities of the Vibrational Transitions of Molecular Hydrogen Ion ( $H_2^+$ ) in the Magnetic Field of a Neutron Star", Astron. Zh. 60(1), 105-9 (1983).
41. Samsonov, B. F., "Calculation of Diatomic Molecules by the MO LCAO Method.  $1s\sigma_g$  and  $2p\sigma_u$  States of  $H_2^+$  and  $1s\sigma$  State of  $HeH_2^+$ ", Izv. Vyssh. Uchebn. Zaved., Fiz. 25(10), 117-19 (1982).
42. Sataka, Masao; Shirai, Toshizo; Kikuchi, Akira; Nakai, Yohta, "Ionization Cross Sections for Ion-Atom and Ion-Molecule Collisions. I. Ionization Cross Sections for Hydrogen and Helium Ions  $H^+$ ,  $H_2^+$ ,  $H_3^+$ ,  $He^+$  and  $He^{++}$  Incident on H,  $H_2$ , and He", Nippon Genshiryoku Kenkyusho, [Rep.] JAERI-M, JAERI-M-9310, 65 Pp. (1981).
43. Cue, N.; Edwards, A. K.; Gemmell, D. S.; Kanter, E. P.; Kutina, R., "UV-Laser Photofragmentation of a 2-MeV Molecular Hydrogen Ion ( $H_2^+$ ) Beam", Ann. Isr. Phys. Soc., 4 (Mol. Ions, Mol. Struct. Interact. Matter), 194-7 (1981).
44. Sonnleitner, Stephanie A.; Beckel, Charles L.; Colucci, Anthony J.; Scaggs, E. Rodney, "Rational Fraction Representation of Diatomic Vibrational Potentials. V. The  $3d\sigma_g$  State of  $H_2^+$ ", J. Chem. Phys. 75(4), 2018-20 (1981).



45. Anderson, S. L.; Hirooka, T.; Tiedemann, P. W.; Mahan, B. H.; Lee, Y. T., "Photoionization of Hydrogen  $[(H_2)_2]$  and Clusters of Oxygen Molecules", J. Chem. Phys. 73(10), 4779-83 (1980).
46. Huber, B. A.; Schulz, U.; Wiesemann, K., "Cross Sections for Slow Ion Production and Charge Transfer in  $H_3^+(D_3^+)-H_2(D_2)$  Collisions", Phys. Lett. A, 79A(1), 58-60 (1980).
47. Hirooka, Tomohiko; Anderson, Scott L.; Tiedemann, Peter W.; Mahan, Bruce H.; Lee, Yuan T., "Photoionization of Molecular Hydrogen Dimer  $(H_2)_2$ ", Koen Yoshishu - Bunshi Kozo Sogo Toronkai, 64-5. Chem. Soc. Japan: Tokyo, Japan. (1979).
48. Kaschiev, M. S.; Vinitskii, S. I.; Vukajlovic, F. R., "Hydrogen Atom H and Hydrogen  $(H_2^+)$  Molecule in Strong Magnetic Fields", Phys. Rev. A, 22(2), 557-9 (1980).
49. Nir, D.; Navon, E.; Mann, A.; Rosner, B., "Relations Between the Dissociation Cross Section of  $H_2^+$  Ions and the Charge Exchange Cross Sections of Their Fragments", Phys. Lett. A, 77A(2-3), 150-2 (1980).
50. Koyano, Inosuke; Tanaka, Kenichiro, "State-Selected Ion-Molecule Reactions by a Threshold Electron-Secondary Ion Coincidence (TESICO) Technique. I. Apparatus and the Reaction  $H_2^+ + H_2 \rightarrow H_3^+ + H$ ", J. Chem. Phys. 72(9), 4858-68 (1980).
51. Semenov, V. E., "Calculation of the Cross Section for Photoionization of a Molecular Hydrogen Ion  $H_2^+$  Based on Gaussian Functions", Opt. Spektrosk. 48(4), 723-7 (1980).
52. Oda, Nobuo; Urakawa, Junji; Tokoro, Nobuhiro; Nojiri, Hiroshi, "Ionizing Collisions of Hydrogen Cluster Ions ( $H^+$ ,  $H_2^+$ ,  $H_3^+$ ) of Intermediate Energies with Gases", Iongen to Sono Oyo, Shinpojumu, 4th, 205-8. Ion Kogaku Kondankai: Kyoto, Japan. (1980).
53. Dastidar, K. Rai; Bose, M.; Dastidar, T. K. Rai, "Electron Cooling Through Resonant Collisions with Hydrogen  $(H_2^+)$  Molecular Ion", J. Phys. Soc. Jpn. 47(6), 1955-8 (1979).
54. Boikova, R. F., "Ion-Ion Recombination of Molecular Hydrogen Ion  $(H_2^+)$  and Hydrogen Ion  $(H^+)$ ", Vestn. Leningr. Univ., Fiz., Khim. 2, 103-5 (1979).
55. Moiseyev, Nimrod; Corcoran, Chris, "Autoionizing States of Diatomic Hydrogen and Hydrogen Ion  $(H_2^+)$  Using The Complex-Scaling Method", Phys. Rev. A, 20(3), 814-17 (1979).
56. Propin, R. Kh., "Exponentially Small Part of the Molecular Hydrogen  $(H_2^+)$  Ion Wave Function", Latv. PSR Zinat. Akad. Vestis, Fiz. Teh. Zinat. Ser. 3, 7-12 (1979).

57. Hashemi-Attar, Ali-Reza; Beckel, Charles L.; Keepin, William N.; Sonnlleitner, Stephanie A., "A New Functional Form for Representing Vibrational Eigenenergies of Diatomic Molecules. Application to Hydrogen Molecular Ion ( $H_2^+$ ) Ground State", J. Chem. Phys. 70(8), 3881-3 (1979).
58. Strand, Michael P.; Reinhardt, William P., "Semiclassical Quantization of the Low-Lying Electronic States of Hydrogen Molecular Ion ( $H_2^+$ )", J. Chem. Phys. 70(8), 3812-27 (1979).
59. Bishop, David M.; Cheung, Lap M., "Natural Orbital Analysis of Nonadiabatic  $H_2^+$  Wave Functions", Int. J. Quantum Chem. 15(5), 517-32 (1979).
60. Dastidar, K. Rai; Dastidar, T. K. Rai, "Dissociative Recombination of  $H_2^+$ ,  $HD^+$ , and  $D_2^+$  Molecular Ions", J. Phys. Soc. Jpn. 46(4), 1288-94 (1979).
61. Van der Hart, Johanna A.; Mulder, J. J. C., "Ab Initio CI LCAO Calculations of the Lowest Core-Excited  $\Sigma$  Resonant States of  $H_2^-$ ", Chem. Phys. Lett. 61(1), 111-14 (1979).
62. Beck, D. R.; Nicolaides, C. A., "How Many Bound States do Hydrides  $H^-$  and  $H_2^-$  Have?", Chem. Phys. Lett. 59(3), 525-8 (1978).
63. Teloy, E., "Proton-Hydrogen ( $H_2$ ,  $D_2$ ) Differential Inelastic and Reactive Scattering At Low Energies", Electron. At. Collisions, Proc. Int. Conf., 10th. Meeting Date 1977, 591-603. Edited by: Watel, Guy. North-Holland: Amsterdam, Neth. (1978).
64. Veselov, M. G.; Rekasheva, T. N., "Extension of the LCAO Method [Hydrogen ( $H_2^+$ ) Ion Calculation]", Vestn. Leningr. Univ., Fiz., Khim. (2), 14-16 (1978).
65. Tame, B. R.; Ritchie, Burke, "Continuum States for Hydrogen Molecule: A Study of Convergence in  $e^-H_2^+$  Scattering Equations", J. Chem. Phys. 68(8), 3595-9 (1978).
66. Stine, J. R.; Muckerman, J. T., "Charge Exchange and Chemical Reaction in the  $H_2^+ + H_2$  System. I. Characterization of the Potential Energy Surfaces and Nonadiabatic Regions", J. Chem. Phys. 68(1), 185-94 (1978).
67. Gentry, W. Ronald; Ringer, Geoffrey, "On the Possibility That Electronically Excited Products May Be Formed in the Reaction  $H_2^+ + H_2 \rightarrow H_3^+ + H$ ", J. Chem. Phys. 67(11), 5398-9 (1977).
68. Douglass, Charles H.; McClure, Donald J.; Gentry, W. Ronald, "The Dynamics of the Reaction  $H_2^+ + H_2 \rightarrow H_3^+ + H$ , with Isotopic Variations", J. Chem. Phys. 67(11), 4931-40 (1977).
69. Malkhasyan, R. T.; Zhurkin, E. S.; Tunitskii, N. N., "Excitation of Hydrogen Ions ( $H_3^+$ ) and Dependence of the Cross Section of the Secondary

Ionic-Molecular Reaction  $H_3^+ + Ar \rightarrow ArH^+ + H_2$  on Excitation Energy of the  $H_3^+$  Ion", Khim. Vys. Energ. 11(6), 400-2 (1977).

70. Top, Zvi H.; Baer, Michael, "Incorporation of Electronically Nonadiabatic Effects Into Bimolecular Reactive Systems. II. The Collinear ( $H_2 + H^+$ ,  $H_2^+ + H$ ) System", Chem. Phys. 25(1), 1-18 (1977).
71. Steinborn, E. Otto; Weniger, E. J., "Advantages of Reduced Bessel Functions as Atomic Orbitals. An Application to the Hydrogen Ion ( $H_2^+$ )", Int. J. Quantum Chem., Symp., 11 (Proc. Int. Symp. At., Mol., Solid-State Theory, Collision Phenom., Comput. Methods), 509-16 (1977).
72. Rekasheva, T. N., "Calculation of the Hydrogen ( $H_2^+$ ) Molecular Ion", Tr. Leningr. Korablestroit. Inst. 97, 60-1 (1975).
73. Hyatt, D.; Careless, P. N.; Stanton, L., "A Simple Ab-initio Potential Surface for the Reaction  $H_3^+(H_2, H_2)H_3^+$  in  $C_{2v}$  Symmetry", Int. J. Mass Spectrom. Ion Phys. 23(1), 45-50 (1977).
74. Auerbach, D.; Cacak, R.; Caudano, R.; Gaily, T. D.; Deyser, C. J.; McGowan, J. Wm.; Mitchell, J. B. A.; Wilk, S. F. J., "Merged Electron-Ion Beam Experiments I. Method and Measurements of ( $e-H_2^+$ ) and ( $e-H_3^+$ ) Dissociative-Recombination Cross Sections", J. Phys. B: Atom. and Molec. Phys. 10(18), 3797-820 (1977).
75. Alvarez, Ignacio; Cisneros, Carmen; Barnett, C. F.; Ray, J. A., "Negative-Ion Formation From Dissociative Collisions of  $H_2^+$ ,  $H_3^+$ , and  $HD_2^+$  in Molecular Hydrogen, Helium, and Xenon", Phys. Rev. A, 14(2), 602-7 (1976).
76. Adamov, M. N.; Ivanov, A. I., "Calculation of Electric Field Gradient in Hydrogen Ion ( $H_2^+$ ) on a Four-Center Base From Slater Type Functions", Vestn. Leningr. Univ. Fiz., Khim. (1), 28-31 (1976).
77. Vestal, M. L.; Blakley, C. R.; Ryan, P. W.; Futrell, J. H., "Crossed-Beam Study of the Reaction  $H_3^+(D_2, H_2)D_2H^+$ ", J. Chem. Phys. 64(5), 2094-111 (1976).
78. Ozenne, J.-B.; Durup, J.; Odom, R. W.; Pernot, C.; Tabché-Fouhaillé, A.; Tadjeddine, M., "Laser Photodissociation of the Isotopic Hydrogen Molecular Ions. Comparison Between Experimental and Ab Initio Computed Fragment Kinetic Energy Spectra", Chem. Phys. 16, 75-80 (1976).
79. Barsuk, A. A.; Zenchenko, V. P.; Isanov, M. M., "Hydrogen Molecular Ion  $H_2^+$  in a Magnetic Field", Tezisy Dokl. Soobshch.-Konf. Molodykh Uch. Mold., 9th, Meeting Date 1974, 50. Edited by: Lazarev, A. M. "Shtiintsa": Kishinev, USSR. (1975).
80. Lees, A. B.; Rol, P. K., "Merging Beams Study of the  $D^+(H_2, H)HD^+$  and

$H_2^+(D,H)HD^+$  Reaction Mechanisms", J. Chem. Phys. 63(6), 2461-5 (1975).

81. Bishop, David M.; Shih, Shing-Kuo; Beckel, Charles L.; Wu, Fun-Min; Peek, James M., "Theoretical Study of  $H_2^+$  Spectroscopic Properties. IV. Adiabatic Effects for the  $2p\pi_u$  and  $3d\sigma_g$  Electronic States", J. Chem. Phys. 63(11), 4836-41 (1975).
82. Elford, M. T.; Milloy, H. B., "Mobility of  $H_3^+$  and  $H_5^+$  Ions in Hydrogen and the Equilibrium Constant for the Reaction  $H_3^+ + 2H_2 = H_5^+ + H_2$  at Gas Temperatures of 195, 273, and 293 K", Aust. J. Phys. 27(6), 795-811 (1974).
83. Hiraoka, K.; Kebarle, P., "Temperature Dependence of Third Order Ion Molecule Reactions. Reaction  $H_3^+ + 2H_2 = H_5^+ + H_2$ ", J. Chem. Phys. 63(2), 746-9 (1975).
84. Hiraoka, K.; Kebarle, P., "Determination of the Stabilities of  $H_5^+$ ,  $H_7^+$ ,  $H_9^+$ , and  $H_{11}^+$  From Measurement of the Gas Phase Ion Equilibria  $H_n^+ + H_2 = H_{n+2}^+$  ( $n = 3, 5, 7, 9$ )", J. Chem. Phys. 62(6), 2267-70 (1975).
85. Robiette, Alan G., "Variation Theorem Applied to Hydrogen Ion ( $H_2^+$ ). Simple Quantum Chemistry Computer Project", J. Chem. Educ. 52(2), 95-6 (1975).
86. Lees, A. E.; Rol, P. K., "Merging Beams Study of the  $H_2^+(H_2,H)H_3^+$ ,  $H_2^+(D_2,H)HD_2^+$ , and  $D_2^+(H_2,H)HD_2^+$  Reaction Mechanisms", J. Chem. Phys. 61(11), 4444-9 (1974).
87. Wendell, K. L.; Rol, P. K., "Merging Beams Study of the Reaction  $H_2^+(D,H)HD^+$ ", J. Chem. Phys. 61(5), 2059-61 (1974).
88. Malkhasyan, R. T.; Zhurkin, E. S.; Tikhomirov, M. V.; Tunitskii, N. N., "Charge Exchange of Argon(1+) and Hydrogen( $H_2^+$ ) Ions with Deuterium Molecules in Reactive Collisions At Up to 100 eV, and Formation of Deuterium ( $D_3^+$ ) Ions in Ion-Molecule Reactive Collisions Involving the Charge-Exchanged Deuterium ( $D_2^+$ )<sup>+</sup> System", Chem. Phys. Lett. 119(2-3), 169-72 (1985).
89. Van Assalt, N. P. F. B.; Maas, J. G.; Los, J., "Laser Induced Photodissociation of  $H_2^+$  and  $D_2^+$  Ions", Chem. Phys. 5, 429-38 (1974).
90. Peart, B; Dolder, K. T., "Collisions Between Electrons and  $H_2^+$  Ions. V. Measurements of Cross Sections for Dissociative Recombination", J. Phys. B: Atom. Molec. Phys. 7(2), 236-43 (1974).
91. Davydkin, V. A.; Rapoport, L. P., "Two-Photon Ionization of Molecular Hydrogen Ion ( $H_2^+$ )", Opt. Spektrosk. 36(2), 244-9 (1974).
92. Bunker, P. R., "Nonadiabatic Effects on the Vibrational Intervals of Hydrogen Molecular Ion [ $H_2^+$ ]", J. Mol. Spectrosc. 46(3), 504-5 (1973).

93. Zhurkin, E. S.; Kaminskii, V. A.; Tikhomirov, M. V.; Tunitskii, N. N., "Cross Sections of Hydrogen ( $H_2^+$ ) and Deuterium ( $D_2^+$ ) Ion Dissociation in the 0.1-2 KeV Range", Zh. Tekh. Fiz. 43(2), 405-9 (1973).
94. Beckel, C.; Shapi, M.; Peek, J. M., "Theoretical Study of  $H_2^+$  Spectroscopic Properties. II. The  $2p\pi_u$  Electronic State", J. Chem. Phys. 59(10), 5288-93 (1973).
95. Stewart, Ronald F., "Finite-Difference Solution of the United Atom, One-Center Expansion for  $H_2^+$ ", Mol. Phys. 24(4), 879-83 (1972).
96. Bochvar, D. A.; Tutkevich, A. V., "Chemical Bonding and Entropy of Electron Density Distribution in Molecules. II. Characteristics of Electron Density Distribution in the  $H_2^+$  Molecular Ion", Zh. Strukt. Khim. 13(4), 678-81 (1972).
97. Herrero, F. A.; Doering, J. P., "Superelastic Collisions of Vibrationally Excited  $H_2^+$  with Atoms and Molecules", Phys. Rev. Lett. 29(10), 609-11 (1972).
98. Sauers, I.; Fitzwilson, R. L.; Ford, J. C.; Thomas, E. W., "Angular Distribution of Metastable Hydrogen Formed by Dissociation of  $H_2^+$ ", Phys. Rev. A, 6(4), 1418-24 (1972).
99. Ali, M. K.; Meath, W. J., "A Floating One-Center Perturbation Treatment for  $H_2^+$ -Like Molecules", Int. J. Quantum Chem. 6(5), 949-66 (1972).
100. Peart, B.; Dolder, K. T., "Collisions Between Electrons and  $H_2^+$  Ions. III. Measurements of Proton Production Cross Sections at Low Energies", J. Phys. B, 5(8), 1554-8 (1972).
101. Tung, Eleanor W.; Sanders, William A., "Simple Perturbation and Perturbation-Variation Treatments of the  $1s\sigma_g$  and  $2p\sigma_u$  States of  $H_2^+$ ", Int. J. Quantum Chem. 6(4), 717-23 (1972).
102. Dolder, K.; Peart, B., "Proton Production by Collisions Between Electrons and  $H_2^+$ . Comments", J. Phys. B, 5(6), L129-L131 (1972).
103. Taylor, Howard S.; Thomas, Lowell D., "Short-Lived Resonant State of  $H_2^-$ ", Phys. Rev. Lett. 28(17), 1091-2 (1972).
104. Rundel, R. D., "Proton Production in Collisions Between Electrons and  $H_2^+$  Ions", J. Phys. B, 5(4), L76-L78 (1972).
105. Weinhold, Frank; Chinen, Allan B., "Variational Wave Functions for  $H_2^+$ ", J. Chem. Phys. 56(8), 3798-801 (1972).
106. Von Busch, Friedrich; Dunn, Gordon H., "Photodissociation of  $H_2^+$  and  $D_2^+$ : Experiment", Phys. Rev. A 5(4), 1726-43 (1972).

107. Ozenne, J.-B.; Pham, D.; Durup, J., "Photodissociation of  $H_2^+$  by Monochromatic Light with Energy Analysis of the Ejected  $H^+$  Ions", Chem. Phys. Lett. 17(3), 422-4 (1972).
108. Kelkar, V. K.; Bhalla, K. C.; Khubchandani, P. G., "Study of the Hydrogen Molecule Using  $H_2^+$  Molecular Orbital", Mol. Phys. 22(6), 1141-3 (1971).
109. Borkowski, Jozef, "Calculations of the Photoionization Cross Section of  $H_2^+$  from  $1s\sigma_g$  State by Using Some Approximate Wave Functions", Bull. Soc. Sci. Lett. Lodz, 24(3), 4 Pp. (1970).
110. Peart, B.; Dolder, K. T., "Collisions Between Electrons and  $H_2^+$  Ions. I. Measurements of Cross Sections for Proton Production", J. Phys. B, 4(11), 1496-505 (1971).
111. Laurenzi, Bernard J., "Green's Function for the Hydrogen Molecular Ion  $H_2^+$ ", J. Chem. Phys. 55(6), 2681-4 (1971).
112. Jackson, M.; McEachran, R. P.; Cohen, M., "James Wave Function for the Ground State of  $H_2^+$ ", Chem. Phys. Lett. 10(2), 143-5 (1971).
113. Borkman, R. F., "Electric Quadrupole Moments for  $H_2^+$ ,  $H_2$ , and  $H_3^+$  from a Point-Charge Model", Chem. Phys. Lett. 9(6), 624-6 (1971).
114. Borisov, M. S.; Vetchinkin, S. I., "Threshold Photoionization of the  $H_2^+$  Molecular Ion in the United Atom Model", Opt. Spektrosk. 30(3), 409-12 (1971).
115. Kawaoka, Kenji; Borkman, Raymond F., "Single-Center Calculations on the Electronically Excited States of Equilateral  $H_3^+$  Ion", J. Chem. Phys. 54(10), 4234-8 (1971).
116. Trivedi, P. C., "Wave Functions for  $H_2^+$ ", J. Phys. B, 4(4), 420-3 (1971).
117. Bishop, David M., "Cubic and Quartic Force Constants for  $H_2^+$ ", J. Chem. Phys. 54(6), 2761-2 (1971).
118. Katriel, J.; Adam, Gabriel, "Exact Analytic Evaluation of the  $H_2^+$  Force Constant", Chem. Phys. Lett. 8(2), 191-4 (1971).
119. Mueller, Hans, "Theoretical Study of Hydrogen Polymers. I. Calculation of the Proton Affinities of H,  $H_2$ , and  $H_3^+$ ", Z. Chem. 10(12), 478-9 (1970).
120. Silters, E.; Ustinov, N. N.; Yurkevich, V. E.; Bolotin, A. B., "Calculation of the Energy Spectrum and Wave Functions of the  $H_2^+$  Ion Electron", Str. Mol. Kvantovaya Khim., 138-42. Edited by: Brodskii, A. I. "Naukova Dumka": Kiev, USSR. (1970).

121. Jasperse, J. R., "Method for One Particle Bound to Two Identical Fixed Centers: Application to  $H_2^+$ ". Phys. Rev. A, [3]2(6), 2232-44 (1970).
122. Tunitskii, N. N.; Zhurkin, E. S.; Tikhomirov, M. V., "Effect of the Excitation of  $H_2^+$  and  $D_2^+$  Ions on the Cross Section of Their Dissociation During a Collision with Atoms and Molecules At 0.1-2 keV", Pis'ma Zh. Eksp. Teor. Fiz. 12(6), 312-14 (1970).
123. Weingartshofer, A.; Ehrhardt, Helmut; Hermann, V.; Linde, F., "Measurements of Absolute Cross Sections for (e,  $H_2$ ) Collision Processes. Formation and Decay of  $H_2^-$  Resonances", Phys. Rev. A, [3] 2(2), 294-304 (1970).
124. Schaad, Lawrence J.; Hicks, W. V., "Equilibrium Bond Length in  $H_2^+$ ", J. Chem. Phys. 53(2), 851-2 (1970).
125. Chang, Edward S.; Temkin, Aaron, "Rotational Excitation of Diatomic Molecular Systems. II.  $H_2^+$ ", J. Phys. Soc. Jap. 29(1), 172-9 (1970).
126. Csizmadia, Imre G.; Kari, R. E.; Polanyi, John C.; Roach, A. C.; Fobb, M. A., "Ab-initio SCF-MO-CI Calculations for  $H^-$ ,  $H_2$ , and  $H_3^+$  Using Gaussian Basis Sets", J. Chem. Phys. 52(12), 6205-11 (1970).
127. Burt, J. A.; Dunn, Jerry L.; McEwan, M. J.; Sutton, M. M.; Roche, A. E.; Schiff, Harold I., "Ion-Molecule Reactions of  $H_3^+$  and the Proton Affinity of  $H_2$ ", J. Chem. Phys. 52(12), 6062-75 (1970).
128. Schaad, L. J.; Hicks, W. V., "Equilibrium Bond Length in  $H_2^+$ ", J. Chem. Phys. 53(2), 851-52 (1970).
129. Beckel, Charles L.; Hansen, Bertle D. III, "Theoretical Study of  $H_2^+$  Ground Electronic State Spectroscopic Properties", J. Chem. Phys. 53(9), 3681-90 (1970).
130. Jefferts, Keith B., "Hyperfine Structure in the Molecular Ion  $H_2^+$ ", Phys. Rev. Lett. 23(26), 1476-7 (1969).
131. Jackson, Malcolm; McEachran, Robert P.; Cohen, Maurice, "Second-Order Perturbation Treatment of the Ground State of  $H_2^+$ ", J. Chem. Phys. 52(1), 102-6 (1970).
132. Calvert, J. McI.; Davison, William Donald, "Numerical Single-Center Calculation of the Polarizabilities of  $H_2^+$ ", Chem. Phys. Lett. 4(6), 327-30 (1969).
133. Bishop, David M.; Macias, A., "Ab Initio Calculation of Harmonic Force Constants. II. Application to Gaussian Wavefunctions for  $H_2^+$ ", J. Chem. Phys. 51(11), 4997-5001 (1969).

134. Temkin, Aaron; Vasavada, K. V.; Chang, Edward S.; Silver, A., "Scattering of Electrons From  $H_2^+$ . II", Phys. Rev. 186(1), 57-66 (1969).
135. Stecher, Theodore P.; Williams, David A., "Interstellar  $H_2^+$  Molecule", Astrophys. Lett. 4(3), 99-101 (1969).
136. Lyon, William D.; Matcha, Robert L.; Sanders, William A.; Meath, William J.; Hirschfelder, Joseph O., "Erratum: Perturbation Treatment of the Ground State of  $H_2^+$ ", J. Chem. Phys. 51(7), 3151-2 (1969).
137. Latimer, C. J.; Browning, R.; Gilbody, H. B., "Dissociation and Charge Transfer in 1.4-46 keV  $H_2^+-H_2$  Collisions", Proc. Phys. Soc., London (At. Mol. Phys.), [2]2(10), 1055-9 (1969).
138. Bochvar, D. A.; Borodzich, R. M.; Tutkevich, A. V., "Chemical Bonding and Entropy of Electron Density Distribution in Molecules. I.  $H_2^+$  and  $H_2$ ", Zh. Strukt. Khim. 10(3), 530-2 (1969).
139. Brandas, Erkki; Goscinski, O., "Symmetry-Adapted Second-Order Energy. Some Comments and Results for  $H_2^+$ ", J. Chem. Phys. 51(3), 975-83 (1969).
140. Meierjohann, B.; Seibt, W., "Collision-Induced Dissociation of  $H_2^+$  Ions Energy and Angular Dependence", Z. Phys. 225(1), 9-25 (1969).
141. Luke, S. K., "Radiofrequency Spectrum of  $H_2^+$ ", Astrophys. J. 156(2)(Pt. 1), 761-9 (1969).
142. Radel, Stanley R.; Gorman, Ronnie; Cutler, Carol; Kahn, Luis, "Geometric-Mean Variation Function for the Hydrogen Molecular Ion  $H_2^+$ ", J. Chem. Phys. 50(8), 3642-4 (1969).
143. Leventhal, Jacob J.; Friedman, Lewis, "Energy Transfer in the De-Excitation of  $H_3^+$  by  $H_2$ ", J. Chem. Phys. 50(7), 2928-31 (1969).
144. Conroy, Harold, "Molecular Schrödinger Equation. X. Potential Surfaces for Ground and Excited States of Isosceles  $H_3^{++}$  and  $H_3^+$ ", J. Chem. Phys. 51(9), 3979-3993 (1969).
145. Luke, S. K.; Hunter, G.; McEachran, Robert P.; Cohen, Maurice, "Relativistic Theory of  $H_2^+$ ", J. Chem. Phys. 50(4), 1644-50 (1969).
146. Peek, James M., "Discrete Vibrational States Due Only to Long-Range Forces:  $^2\Sigma_u^+$  ( $2p\sigma_u$ ) State of  $H_2^+$ ", J. Chem. Phys. 50(10), 4595-96 (1969).
147. Luke, S. K.; Hunter, G.; McEachran, R. P.; Cohen, M., "Relativistic Theory of  $H_2^+$ ", J. Chem. Phys. 50(4), 1644-50 (1969).
148. Gersten, Joel I., "Evaluation of the Lamb Shift for the Hydrogen Molecule-Ion", J. Chem. Phys. 51(8), 3181-85 (1969).



149. Dunn, Gordon H., "Photodissociation of  $H_2^+$  and  $D_2^+$ : Theory and Tables", Joint Inst. Lab. Astrophys., Rep. No. 92, 55 Pp. (1968).
150. Tunitskii, N. N., "Effect of the Excitation of  $H_2^+$  And  $D_2^+$  Ions in the Reaction in Which They Take Part", Teor. Eksp. Khim. 4(5), 695-8 (1968).
151. Winter, Nicholas Wilhelm; McKoy, Vincent, "Numerical One-Center Calculation of the ns-( $\sigma$ ) Rydberg Series of  $H_2^+$ ", J. Chem. Phys. 49(10), 4728-30 (1968).
152. Bishop, David M., "Calculation of the Electric Field Gradient at the Nucleus in  $H_2^+$ ", J. Chem. Phys. 49(8), 3718-22 (1968).
153. Burke, P. G., "Potential Energy Curves of  $H_2^-$ ", Proc. Phys. Soc., London, At. Mol. Phys. [2]1(4), 586-8 (1968).
154. Main, I. G.; Durell, J. L.; Sareen, R. A., "Search for the Formation of  $H_2^-$  Ions From 40-keV  $H_2^+$  Ions in Hydrogen", Proc. Phys. Soc., London, At. Mol. Phys. [2] 1(4), 755-7 (1968).
155. Hayden, Howard C.; Amme, Robert C., "Vibrational Excitation Effects on Charge-Transfer Processes Involving  $H_2^+$  and  $D_2^+$  Between 70 and 1000 eV", Phys. Rev. 172(1), 104-9 (1968).
156. Damburgs, R.; Propin, R. Kh., "Asymptotic Expansion of the Electronic Terms of the Hydrogen Molecular Ion  $H_2^+$ ", Latv. PSR Zinat. Akad. Vestis, Fiz. Teh. Zinat. Ser. (1), 50-9 (1968).
157. Jefferts, Keith B., "Rotational Hyperfine Structure Spectra of Hydrogen Molecular Ions", Phys. Rev. Lett. 20(2), 39-41 (1968).
158. Blicharski, Jerzy S., "Radio-Frequency Transitions and Nuclear and Electron Polarization for Hydrogen Molecular Ions  $H_2^+$ ", Can. J. Phys. 46(7), 823-9 (1968).
159. Hughes, Raymond Hargett; Kay, David B.; Stigers, C. A.; Stokes, E. D., "Production of Hydrogen Atoms in the 3s State by the Dissociation of Fast  $H_2^+$  and  $H_3^+$  Projectiles on Impact with Hydrogen, Helium, Argon, and Neon", Phys. Rev. 167(1), 26-9 (1968).
160. Neynaber, Roy H.; Trujillo, S. M., "Study of  $H_2^+ + H_2 \rightarrow H_3^+ + H$  Using Merging Beams", Phys. Rev. 167(1), 63-6 (1968).
161. Richardson, C. B.; Jefferts, K. B.; Dehmelt, H. G., "Alignment of the  $H_2^+$  Molecular Ion by Selective Photodissociation. II. Experiments on the Radio-Frequency Spectrum", Phys. Rev. 165(1), 80-87 (1968).
162. Solov'ev, E. S.; Il'in, R. N.; Oparin, V. A.; Fedorenko, N. V., "Produc-

tion of Highly Excited Hydrogen Molecules and Atoms by Fast  $H_2^+$  and  $H_3^+$  Ions Passing Through Hydrogen, Neon, and Magnesium and Sodium Vapor", Zh. Eksp. Teor. Fiz. 53(6), 1933-41 (1967).

163. Oksyuk, Yu. D., "Effect of Vibrational Excitation on the Dissociation of the Molecular Hydrogen Ion Under Electron and Proton Bombardment", Opt. Spektrosk. 23(3), 366-73 (1967).
164. Williams James Francis; Dunbar, D. N. F., "Charge Exchange and Dissociation Cross Sections for  $H_1^+$ ,  $H_2^+$ , and  $H_3^+$  Ions of 2- to 50-keV Energy Incident Upon Hydrogen and the Inert Gases", Phys. Rev. 149(1), 62-9 (1966).
165. Bhalla, K. C.; Khubchandani, P. G., "A Variational Function for the Ground State of  $H_2^+$ ", Proc. Phys. Soc., London, 92(3), 529-30 (1967).
166. Dance, D. F.; Harrison, M. F. A.; Rundel, Robert D.; Smith, A. C. H., "A Measurement of the Cross Section for Proton Production in Collisions Between Electrons and  $H_2^+$  Ions", Proc. Phys. Soc., London, 92(3), 577-8 (1967).
167. Eliezer, Isaac; Taylor, Howard S.; Williams, James Kendree, Jr., "Resonant States of  $H_2^-$ ", J. Chem. Phys. 47(6), 2165-77 (1967).
168. McQuarrie, Donald A.; Hirschfelder, Joseph O., "Intermediate-Range Intermolecular Forces in  $H_2^+$ ", J. Chem. Phys. 47(5), 1775-80 (1967).
169. Cohen, Maurice; McEachran, Robert P.; McPhee, Sheila D., "Approximate Molecular Orbitals. IV. The  $3d_g$  and  $4f_u$  States of  $H_2^+$ ", Can. J. Phys. 45(8), 2533-42 (1967).
170. Carlson, Charles M., "Use of Linear Nonsymmetric Eigenvalue Equations for a Sum-Over-Points Approach: Application to  $H_2^+$ ", J. Chem. Phys. 47(2), 862-3 (1967).
171. Silbey, Robert, "Perturbation of Calculation of the Energy of the First Excited State ( $2p\sigma$ ) of  $H_2^+$ ", J. Chem. Phys. 46(10), 4026-8 (1967).
172. Komarov, I. V.; Slavyanov, S. Yu., "Wavefunctions and Electron Terms of the  $H_2^+$  Molecular Ion for Large Internuclear Distances", Zh. Eksp. Teor. Fiz. 52(5), 1368-77 (1967).
173. Patel, Jashbhai C., "Accurate Wavefunction of  $H_2^+$ ", J. Chem. Phys. 47(2), 770-4 (1967).
174. Dei-Cas, Renato; Fumelli, Michele; Girard, Jean Pierre; Prevot, Francois; Valckx, Franciscus P. G., "Dissociation of Hydrogen Molecular Ions by the Lorentz Force", Nucl. Fusion, 6(3), 212-14 (1966).

175. Peek, James M.; Green, Thomas Allen; Weihofen, W. H., "Theory of High-Energy Inelastic Collisions Between Molecular Systems. Dissociation of  $H_2^+$  with  $H_2$ ", Phys. Rev. 160(1), 117-24 (1967).
176. Zare, Richard N., "Dissociation of  $H_2^+$  by Electron Impact: Calculated Angular Distribution", J. Chem. Phys. 47(1), 204-15 (1967).
177. Henderson, Richard C.; Ebbing, Darrell D., "Calculation of Electric Field Gradients in  $H_2^+$  and  $H_2$  Using Single-Center Wavefunctions", J. Chem. Phys. 47(1), 69-72 (1967).
178. Adamov, M. N.; Rebane, T. K.; Evarestov, R. A., "Polarizability of the  $H_2^+$  Ion", Opt. Spektrosk. 22(5), 709-13 (1967).
179. Doverspike, L. D.; Champion, Roy L., "Ion-Molecule Reactions of  $D_2^+$  with  $D_2$  and  $H_2$ ", J. Chem. Phys. 46(12), 4718-25 (1967).
180. Leventhal, Jacob J.; Moran, Thomas F.; Friedman, Lewis, "Molecular Resonant Charge-Transfer Processes;  $H_2^+-H_2$  and  $N_2^+-N_2$ ", J. Chem. Phys. 46(12), 4666-72 (1967).
181. Pavlik, Philip I.; Blinder, Seymour M., "Relativistic Effects in Chemical Bonding: The  $H_2^+$  Molecule", J. Chem. Phys. 46(7), 2749-51 (1967).
182. Dunken, Helga H.; Gottschlich, Klaus, "Relation Between Axial Electron Density and Bond Energy for Various Approximation Functions in  $H_2^+$ ", Z. Phys. Chem. (Leipzig), 233(3-4), 231-6 (1966).
183. Peek, James M., "Theory of Dissociation of  $H_2^+$  by Fast Electrons", Phys. Rev. 154(1), 52-6 (1967).
184. Dunn, Gordon H.; Van Zyl, Bert, "Electron Impact Dissociation of  $H_2^+$ ", Phys. Rev. 154(1), 40-51 (1967).
185. Bottiglionni, Franco; Coutant, Jacques; Gadda, Erio, "Dissociation of  $H_2^+$  and  $H_3^+$  in a Lithium Plasma", J. Phys. (Orsay, Fr.), 27(9-10), 599-604 (1966).
186. McClure, Gordon W., "Dissociation of  $H_2^+$  Ions in Collision with Hydrogen Atoms: 3 to 115 keV", Phys. Rev. 153(1), 182-3 (1967).
187. Lefaivre, Jean, "Energy of  $H_2^+$  Ions by the L.C.A.O. Method", Can. J. Phys. 45(1), 228-230 (1967).
188. Hunter, G.; Pritchard, H. O., "Born-Oppenheimer Separation for Three-Particle Systems. III. Applications", J. Chem. Phys. 46(6), 2153-58 (1967).
189. Buchheit, K.; Henkes, W., "Untersuchung der Massenverteilung von Was-

serstoff-Agglomerat-Ionen in Einem Massenspektrometer mit Energiezerlegung", Z Angew. Phys. 24(4), 191-6 (1967).

190. Dutton, J.; Llewellyn-Jones, Frank; Rees, Walter D.; Williams, Edward Malcolm, "Motion of Slow Positive Ions in Gases. IV. Drift and Diffusion of Ions in Hydrogen", Philos. Trans. R. Soc. London, Ser. A, 259(1100), 339-54 (1966).
191. Valckx, Franciscus P. G.; Verveer, Philippe, "Vibrational Dissociation and Cascade Dissociation of  $H_2^+$  Ions by Collisions with Gas Molecules", J. Phys. (Orsay, Fr.), 27(7-8), 480-4 (1966).
192. Cohen, Maurice; Dorrell, Brenda H.; McEachran, Robert P., "Approximate Molecular Orbitals. II. The  $2p\pi_u$  and  $3d\pi_g$  States of  $H_2^+$ ", Can. J. Phys. 44(11), 2827-38 (1966).
193. Cohen, Maurice; McEachran, Robert P., "Approximate Molecular Orbitals. I. The  $1s\sigma_g$  and,  $2p\sigma_u$  States of  $H_2^+$ ", Can. J. Phys. 44(11), 2809-25 (1966).
194. Nielsen, Svend E.; Dahler, John S., "Theory of the Dissociative Recombination and Associative Ionization of Hydrogen", J. Chem. Phys. 45(11), 4060-79 (1966).

Section II:  $H_3^+$  (and  $H_3^-$ )

1. Le Coz, Georges; Tuffin, Firmin, "A Study of a Collision Process Creating Molecular Ions of Hydrogen ( $H_3^+$ ) and Deuterium ( $D_3^+$ )", C. R. Acad. Sci. Ser. 2, 304(8), 295-300 (1987).
2. Schlier, Christoph G.; Vix, Ulrike, "Complex Formation in Proton-Hydrogen Collisions. II. Isotope Effects", Chem. Phys. 113, 211-221 (1987).
3. Mitchell, J. B. A., "The Role of Electron-Ion Recombination in Bulk Antimatter Production", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 143-156. Edited by John T. Bahns. University of Dayton Research Institute (1987).
4. Bahns, J. T., "Key Problems and Hydrogen Atom Formation", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 219-230. Edited by John T. Bahns. University of Dayton Research Institute (1987).
5. Bahns, J. T.; Sando, K. M.; Tardy, D. C.; Stwalley, W. C., "Proceedings of the Hydrogen Cluster Ion Study Group", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, A1-64. Edited by John T. Bahns. University of Dayton Research Institute (1987).
6. Crofton, M., "Infrared Spectroscopy of  $H_3^+$ ", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 119-124. Edited by John T. Bahns. University of Dayton Research Institute (1987).
7. Bowers, M. T., "Formation and Reactivity of Small Hydrogen Cluster Ions", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 61-72. Edited by John T. Bahns. University of Dayton Research Institute (1987).
8. Saxon, R., "Overview of Hydrogen Clusters", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 27-37. Edited by John T. Bahns. University of Dayton Research Institute (1987).
9. Vojtik, Jan; Spirko, Vladimir; Jensen, Per, "Vibrational Energies of  $H_3^+$  and  $Li_3^+$  Based on Diatomics-in-Molecules Potentials", Collect. Czech. Chem. Commun. 51(10), 2057-62 (1986).
10. Johnson, B. R., "Semiclassical Vibrational Eigenvalues of  $H_3^+$ ,  $D_3^+$ , and  $T_3^+$  by the Adiabatic Switching Method", J. Chem. Phys. 86(3), 1445-50 (1987).
11. Michels, H. H.; Montgomery, J. A. Jr., "The Electronic Structure and Stability of the  $H_3^-$  Anion", Chem. Phys. Lett. 139(6), 535-539 (1987).

12. Hamilton, I., "Vibrational Spacings for  $H_3^+$ ", J. Chem. Phys. 87(1), 774-776 (1987).
13. Tennyson, Jonathan; Sutcliffe, Brian T., "The Infrared Spectrum of  $H_3^+$  and Its Isotopomers. A Challenge to Theory and Experiment", J. Chem. Soc., Faraday Trans. 2, 82(8), 1151-62 (1986).
14. Child, M. S., "Semiclassical Method for the Determination of Narrow Multichannel Resonances: Application to Hydrogen Molecular Ion ( $H_3^+$ )", J. Phys. Chem. 90(16), 3595-9 (1986).
15. Pan, Fu Shih; Oka, Takeshi, "Calculated Forbidden Rotational Spectra of the Hydrogen Ion ( $H_3^+$ )", Astrophys. J. 305(1, Pt. 1), 518-25 (1986).
16. Miao, Jingwei; Yang Beifang; Hao, Shizuho; Jiang, Zengxue; Shi, Miangong; Cue, N., "Internuclear Separations From Foil Breakup of Fast  $H_2^+$ ,  $H_3^+$ ,  $D_2^+$  and  $D_3^+$  Molecules", Nucl. Instrum. Methods Phys. Res. Sect. B, B13(1-3), 181-3 (1986).
17. Sen, A.; Mitchell, J. B. A., "Production of Vibrationally Cold Ions Using a Radio-Frequency Storage Ion", Rev. Sci. Instrum. 57(5), 754-756 (1986).
18. Helm, Hanspeter, "Observation of High-n Rydberg Series ( $7 < n < 49$ ) of the  $H_3$  Molecule", Phys. Rev. Lett. 56(1), 42-45 (1986).
19. Foster, S. C.; McKellar, A. R. W.; Peterkin, I. R.; Watson, J. K. G.; Pan, F. S.; Crofton, M. W.; Altman, R. S.; Oka, T., "Observation and Analysis of the  $v_2$  and  $v_3$  Fundamental Bands of the  $H_2D^+$  Ion", J. Chem. Phys. 84(1), 91-99 (1986).
20. J. B. A. Mitchell, "Dissociative Recombination of Molecular Ions", 185-222, Atomic Processes in Electron-Ion and Ion-Ion Collisions, Edited by F. Brouillard, Plenum Publishing Corp. (1986).
21. Sen, Amarjit; Mitchell, J. B. A., "Production of  $H_3^+$  Ions with Low Internal Energy for Studies of Dissociative Recombination", J. Phys. B: At. Mol. Phys. 19, L545-549 (1986).
22. Shpirko, V., "Vibration-Rotation Hamiltonian of  $X_3$  Type Molecules in Internal Coordinates. Vibrational Spectrum of Hydrogen ( $H_3^+$ ) Molecule", Aktual. Probl. Spektroskopii. Materialy Simp. Uchenykh Sots. Stran po Nov. Probl. Spektroskopii, Moskva, 18-22 Uyunya, 1984, M. 26-9 From: Ref. Zh., Fiz. (A-Zh.) 1985, Abstr. No. 8D65 (1985).
23. Adams, N. G.; Smith, D., "Dissociative Recombination of Triatomic Hydrogen Cation ( $H_3^+$ ), Oxomethylum, Nitrogen Hydride ( $N_2H^+$ ) and Monoprotonated Methane ( $CH_5^+$ )", NATO ASI Ser., Ser. C, 157(Mol. Astrophys.), 657-9 (1985).

24. Tennyson, Jonathan; Sutcliffe, Brian T., "A Calculation of the Rovibrational Spectra of the  $H_3^+$ ,  $H_2D^+$  and  $D_2H^+$  Molecules", *Mol. Phys.* 56(5), 1175-83 (1985).
25. Levin, F. S.; Shertzer, J., "Channel-Coupling Array Analysis of Electron Correlation in Hydrogen Molecular Ion ( $H_3^+$ )", *Phys. Rev. A: Gen. Phys.* 32(4), 2062-7 (1985).
26. Phillips, T. G.; Blake, Geoffrey A.; Keene, Jocelyn; Woods, R. Claude; Churchwell, E., "Interstellar Hydrogen Ion ( $H_3^+$ ): Possible Detection of the  $1_{10} \rightarrow 1_{11}$  Transition of  $H_2D^+$ ", *Astrophys. J.* 294(1, Pt. 2), L45-L48 (1985).
27. Amano, T., "Difference-Frequency Laser Spectroscopy of Molecular Ions with a Hollow-Cathode Cell: Extended Analysis of the  $\nu_1$  Band of  $H_2D^+$ ", *J. Opt. Soc. Am. B* 2(5), 790-793 (1985).
28. Watson, J. K. G.; Foster, S. C.; McKellar, A. R. W.; Bernath, P.; Amano, T.; Pan, F. S.; Crofton, M. W.; Altman, R. S.; Oka, T., "The Infrared Spectrum of the  $\nu_2$  Fundamental Band of the Hydrogen ( $H_3^+$ ) Molecular Ion", *Can. J. Phys.* 62(12), 1875-85 (1984).
29. Mitchell, J. B. A.; Ng, C. T.; Forand, L.; Janssen, R.; McGowan, J. William, "Total Cross Sections for the Dissociative Recombination of Hydrogen ( $H_3^+$ ), Hydrogen-Deuterium ( $HD_2^+$ ) and Deuterium ( $D_3^+$ )", *J. Phys. B*, 17(24), L909-L913 (1984).
30. Preiskorn, Aleksandra; Woznicki, Wieslaw, "Variational Calculations for the Ground State of Triatomic Hydrogen Ion ( $H_3^+$ )", *Mol. Phys.* 52(6), 1291-301 (1984).
31. Adams, Nigel G.; Smith, David; Alge, Erich, "Measurements of Dissociative Recombination Coefficients of  $H_3^+$ ,  $HCO^+$ ,  $N_2H^+$ , and  $CH_5^+$  at 95 and 300 K Using the FALP Apparatus", *J. Chem. Phys.* 81(4), 1778-84 (1984).
32. Eaker, Charles W.; Schatz, George C., "Semiclassical Vibrational Eigenvalues of Triatomic Molecules: Application of the FFT Method to Sulfur Dioxide, Water, Hydrogen Ion ( $H_3^+$ ) and Carbon Dioxide", *J. Chem. Phys.* 81(5), 2394-9 (1984).
33. Carrington, Alan; Kennedy, Richard A., "Infrared Photodissociation Spectrum of the Hydrogen ( $H_3^+$ ) Ion", *J. Chem. Phys.* 81(1), 91-112 (1984).
34. MacDonald, Jeffrey A.; Biondi, Manfred A.; Johnsen, Rainer, "Recombination of Electrons with Hydrogen ( $H_3^+$  and  $H_5^+$ ) Ions", *Planet. Space Sci.* 32(5), 651-4 (1984).
35. Burton, P. G.; Von Nagy-Felsobuki, E.; Doherty, G.; Hamilton, M., "Vibration Spectrum of Triatomic Hydrogen Ion ( $H_3^+$ ): A Model Hamilton-

AD-A203-801 CONDENSATION AND STORAGE OF HYDROGEN-CLUSTER IONS (U) 272  
DAYTON UNIV OH RESEARCH INST J T BAHNS NOV 88  
UDR-TR-88-28 AAL-TR-88-056 F04611-81-C-0046

UNCLASSIFIED

F/G 7/2

NL





ian", Chem. Phys. 83(1-2), 83-8 (1984).

36. Bae, Y. K.; Coggiola, M. J.; Peterson, J. R., "Search for  $H_2^-$ ,  $H_3^-$ , and Other Metastable Negative Ions", Phys. Rev. A, 29(5), 2888-90 (1984).
37. Saito, Shuji; Kawaguchi, Kentarou; Hirota, Eizi, "The Microwave Spectrum of the  $H_2D^+$  Ion; The  $2_{20} \leftarrow 2_{21}$  Transition", J. Chem. Phys. 82(1), 45-47 (1985).
38. Okumura, M; Yeh, L. I.; Lee, Y. T., "The Vibrational Predissociation Spectroscopy of Hydrogen Cluster Ions", J. Chem. Phys., 83(7), 3705-6 (1985).
39. Warner, H. E.; Conner, W. T.; Petrnmichl, R. H.; Lemoine, B., "Laboratory Detection of the  $1_{10} \rightarrow 1_{11}$  Submillimeter Wave Transition of the  $H_2D^+$  Ion", J. Chem. Phys. 81(5), 2514 (1984).
40. Bogey, M.; Demuynck, C.; Denis, M.; Destombes, J. L.; Lemoine, B., "Laboratory Measurement of the  $1_{10} - 1_{11}$  Submillimeter Line of  $H_2D^+$ ", Astron. Astrophys. 137(2), L13-L14 (1984).
41. Lubic, Karen G.; Amano, T., "Observation of the  $v_1$  Fundamental Band of  $D_2H^+$ ", Can. J. Phys. 62(12), 1886-1888 (1984).
42. Kutina, R. E.; Edwards, A. K.; Pandolf, R. S.; Berkowitz, J., "UV Laser Photodissociation of Molecular Ions", J. Chem. Phys. 80(9), 4112-9 (1984).
43. Gellene, Gregory I.; Porter, Richard F., "Experimental Observations of Excited Dissociative and Metastable States of Triatomic Hydrogen Radical ( $H_3$ ) in Neutralized Ion Beams", J. Chem. Phys. 79(12), 5975-81 (1983).
44. Oka, Takeshi, "The Hydrogen ( $H_3^+$ ) Ion", Mol. Ions: Spectrosc., Struct. Chem., 73-90. Edited by: Miller, Terry Alan; Bondybey, Vladimir E. North-Holland: Amsterdam, Neth. (1983).
45. Mitchell, J. B. A.; Forand, J. L.; Ng, C. T.; Levac, D. P.; Mitchell, R. E.; Mul, P. M.; Claeys, W.; Sen, A.; McGowan, J. William, "Measurement of the Branching Ratio for the Dissociative Recombination of Triatomic Hydrogen ( $H_3^+$ ) + Electron", Phys. Rev. Lett. 51(10), 885-8 (1983).
46. Gaillard, M. J.; De Pinho, A. G.; Poizat, J. C.; Remillieux, J.; Saoudi, R., "Experimental Study of the Triatomic Hydrogen Molecule Through the Collisional Sequence  $H_3^+ \rightarrow H_3 \rightarrow H_3^+$  Undergone by Fast Beams in Argon", Phys. Rev. A, 28(3), 1267-75 (1983).
47. Yamaguchi, Yukio; Gaw, Jeffrey F.; Schaefer, Henry F., III, "Molecular Clustering about a Positive Ion. Structures, Energetics, and Vibrational Frequencies of the Protonated Hydrogen Clusters  $H_3^+$ ,  $H_5^+$ ,  $H_7^+$ , and

- $H_9^+$ ", J. Chem. Phys. 78(6, Pt. 2), 4074-85 (1983).
48. Montgomery, D. L.; Jaecks, D. H., "Three-Body Dissociation of  $H_3^+$ ", Phys. Rev. Lett. 51(20), 1862-1864 (1983).
  49. Elford, M. T., "The Heat of Dissociation of  $H_5^+$  Derived from Measurements of Ion Mobilities", J. Chem. Phys. 79(12) 5951-5959 (1983).
  50. Hirao, K; Yamabe, S., "Theoretical Study on the Structure and Stability of Hydrogen Ion Clusters  $H_n^+$  and  $H_n^-$  ( $n = 3, 5, 7, 9, 11, 13$ )", Chem. Phys. 80, 237-243 (1983).
  51. Poluyanov, L. V., "One Model of High Single-Electron Excitations in the Hydrogen Ion ( $H_3^+$ )", Zh. Strukt. Khim. 23(1), 16-21 (1982).
  52. Rayez, J. C.; Rayez-Meaume, M. T.; Massa, L. J., "Theoretical Study of the Hydride ( $H_3^-$ ) Cluster", J. Chem. Phys. 75(11), 5393-7 (1981).
  53. Adams, N. G.; Smith, D., "A Laboratory Study of the Reaction  $H_3^+ + HD = H_2D^+ + H_2$ : The Electron Densities and the Temperatures in Interstellar Clouds", Astrophys. J. 248(1, Pt. 1), 373-9 (1981).
  54. Goh, S. C.; Swan, J. B., "Collisional Dissociation and the Binding Energy of Hydrogen Molecular Ion ( $H_3^+$ )", Phys. Rev. A, 24(3), 1624-5 (1981).
  55. Sataka, Masao; Shirai, Toshizo; Kikuchi, Akira; Nakai, Yohta, "Ionization Cross Sections for Ion-Atom and Ion-Molecule Collisions. I. Ionization Cross Sections for Hydrogen and Helium Ions  $H^+$ ,  $H_2^+$ ,  $H_3^+$ ,  $He^+$  and  $He^{++}$  Incident on H,  $H_2$ , and He", Nippon Genshiryoku Kenkyusho, [Rep.] JAERI-M, JAERI-M-9310, 65 Pp. (1981).
  56. Shy, J.-T.; Farley, John W.; Wing, William H., "Observation of the Infrared Spectrum of the Triatomic Molecular Ion  $H_2D^+$ ", Phys. Rev. A (24(2), 1146-49 (1981).
  57. Carney, Grady D., "Rotation Energies for Deuterated Hydrogen ( $H_3^+$ ) Oscillators in Zero-Point States of Vibration", Chem. Phys. 54(1), 103-7 (1980).
  58. Oka, Takeshi, "Observation of the Infrared Spectrum of  $H_3^+$ ", Phys. Rev. Lett. 45(7), 531-4 (1980).
  59. Carney, G. D.; Porter, R. N., "Ab Initio Prediction of the Rotation-Vibration Spectrum of  $H_3^+$  and  $D_3^+$ ", Phys. Rev. Lett. 45(7), 537-41 (1980).
  60. Kyrala, George A.; Tolliver, David E.; Wing, William H., "Production of  $H_3^+$ ,  $HeH^+$  and  $He_2^+$  Ion Beams Using a Coaxial Electron-Impact Ion Source", Int. J. Mass Spectrom. Ion Phys. 33(4), 367-82 (1980).

61. Anderson, S. L.; Hirooka, T.; Tiedemann, P. W.; Mahan, B. P.; Lee, Y. T., "Photoionization of Hydrogen  $[(H_2)_2]$  and Clusters of Oxygen Molecules", J. Chem. Phys. 73(10), 4779-83 (1980).
62. Huber, B. A.; Schulz, U.; Wiesemann, K., "Cross Sections for Slow Ion Production and Charge Transfer in  $H_3^+(D_3^+)-H_2(D_2)$  Collisions", Phys. Lett. A, 79A(1), 58-60 (1980).
63. Oda, Nobuo; Urakawa, Junji; Tokoro, Nobuhiro; Nojiri, Hiroshi, "Ionizing Collisions of Hydrogen Cluster Ions ( $H^+$ ,  $H_2^+$ ,  $H_3^+$ ) of Intermediate Energies with Gases", Ionen to Sono Oyo, Shinpojumu, 4th, 205-8. Ion Kogaku Kondankai: Kyoto, Japan. (1980).
64. Koyano, Inosuke; Tanaka, Kenichiro, "State-Selected Ion-Molecule Reactions by a Threshold Electron-Secondary Ion Coincidence (TESICO) Technique. I. Apparatus and the Reaction  $H_2^+ + H_2 \rightarrow H_3^+ + H$ ", J. Chem. Phys. 72(9), 4858-68 (1980).
65. Shy, J.-T.; Farley, J. W.; Lamb, Willis E. Jr.; Wing, William H., "Observation of the Infrared Spectrum of the Triatomic Deuterium Molecular Ion  $D_3^+$ ", Phys. Rev. Lett. 45(7), 535-37 (1980).
66. Carney, G. D., "Refinements in the Vibrations Frequencies of  $H_3^+$  and  $D_3^+$ ", Mol. Phys. 39(4) 923-933 (1980).
67. Dyachenko, G. G.; Nemukhin, A. V.; Stepanov, N. F., "Approximation of Potential Surfaces and Solution of a Vibration Problem for the Molecular Hydrogen Ion ( $H_3^+$ )", Vestn. Mosk. Univ., Ser. 2: Khim. 20(5), 416-22 (1979).
68. Dement'ev, A. I.; Simkin, V. Ya., "Calculations of Potential Surfaces of Ground and Excited States of the Hydrogen Ion ( $H_3^+$ ) and the Hydrogen Molecule ( $H_2$ ) by the Method of Configuration Interaction", Deposited Doc., VINITI 1454-78, 48 Pp. Avail. VINITI (1978).
69. Dykstra, Clifford E.; Swope, William C., "The Hydrogen Ion ( $H_3^+$ ) Potential Surface", J. Chem. Phys. 70(1), 1-3 (1979).
70. Hirooka, Tomohiko; Anderson, Scott L.; Tiedemann, Peter W.; Mahan, Bruce H.; Lee, Yuan T., "Photoionization of Molecular Hydrogen Dimer  $(H_2)_2^+$ ", Koen Yoshishu - Bunshi Kozo Sogo Toronkai, 64-5. Chem. Soc. Japan: Tokyo, Japan. (1979).
71. Garcia, R.; Rossi, A.; Russek, A., "Dissociating States of the  $H_3^+$  System", J. Chem. Phys. 70(12), 5463-67 (1979).
72. Bader, Richard F. W.; Nguyen-Dang, T. Tung; Tal, Yoram, "Quantum Topology of Molecular Charge Distributions. II. Molecular Structure and its Change", J. Chem. Phys. 70(9) 4316-29 (1979).

73. Dykstra, Clifford E.; Swope, William C., "The  $H_3^+$  Potential Surface", J. Chem. Phys. 70(1), 1-3 (1979).
74. Saute, Marcel; Laforgue, Alexandre, "Correlation, Ionization, and Attachment Operators Studied as an Example of Triatomic Hydrogen and  $H_3^+$  Complexes in a Diagramatic Method Extended to Open Shell Systems", J. Chim. Phys. Phys.-Chim. Biol. 75(7-8), 679-88 (1978).
75. Gaillard, M. J.; Gemmell, D. S.; Goldring, G.; Levine, I.; Pietsch, W. J.; Poizat, J. C.; Ratkowski, A. J.; Remillieux, J.; Vager, Z.; Zabransky, B. J., "Experimental Determination of the Structure of Hydrogen Ion ( $H_3^+$ )", Phys. Rev. A, 17(6), 1797-803 (1978).
76. Bortsaikin, S. M.; Posashkov, S. A., "Spectral Terms and Wave Functions of a Hydrogen ( $H_2^+$ ) Ion in a Problem of a Hydrogen ( $H_3^+$ ) Ion", Temat. Sb. Nauch. Tr. Mosk. Aviats. in-t, (440), 20-7 (1978).
77. Siegbahn, P.; Liu, B., "An Accurate Three-Dimensional Potential Energy Surface for  $H_3$ ", J. Chem. Phys. 68(5), 2457-65 (1978).
78. Dykstra, Clifford E.; Gaylord, Arthur S.; Gwinn, William D.; Swope, William C.; Schaefer, Henry F. III, "The Uncoupled Symmetric Stretching Frequency of  $H_3^+$ ", J. Chem. Phys. 68(8), 3951-52 (1978).
79. Yamabe, S.; Hirao, K.; Kitaura, K., "Theoretical Study on the Stability and the Structure of  $H_n^+$  ( $n = 3, 5, 7, 9, 11$ )", Chem. Phys. Lett. 56(3), 546-8 (1978).
80. Bardo, Richard D.; Wolfsberg, Max, "The Adiabatic Correction for Nonlinear Triatomic Molecules: Techniques and Calculations", J. Chem. Phys. 68(6) 2686-95 (1978).
81. Lupu, D.; Bucur, R. V., "Properties Suggesting  $H_3^+$ -Type Clusters in Some Metallic Hydrides", J. Phys. Chem. Solids, 38(4), 387-91 (1977).
82. Gentry, W. Ronald; Ringer, Geoffrey, "On the Possibility That Electronically Excited Products May Be Formed in the Reaction  $H_2^+ + H_2 \rightarrow H_3^+ + H$ ", J. Chem. Phys. 67(11), 5398-9 (1977).
83. Douglass, Charles H.; McClure, Donald J.; Gentry, W. Ronald, "The Dynamics of the Reaction  $H_2^+ + H_2 \rightarrow H_3^+ + H$ , with Isotopic Variations", J. Chem. Phys. 67(11), 4931-40 (1977).
84. Malkhasyan, R. T.; Zhurkin, E. S.; Tunitskii, N. N., "Excitation of Hydrogen Ions ( $H_3^+$ ) and Dependence of the Cross Section of the Secondary Ionic-Molecular Reaction  $H_3^+ + Ar \rightarrow ArH^+ + H_2$  on Excitation Energy of the  $H_3^+$  Ion", Khim. Vys. Energ. 11(6), 400-2 (1977).
85. Hyatt, D.; Careless, P. N.; Stanton, L., "A Simple Ab-initio Potential

Surface for the Reaction  $H_3^+(H_2, H_2)H_3^+$  in  $C_{2v}$  Symmetry", Int. J. Mass Spectrom. Ion Phys. 23(1), 45-50 (1977).

86. Alrichs, Reinhart; Votava, Christian; Zirz, Constantin, "Comment: The Bound  $^3\Sigma_u^+$  Excited Level of  $H_3^+$ ", J. Chem. Phys. 66(6), 2771-72 (1977).
87. Auerbach, D.; Cacak, R.; Caudano, R.; Gaily, T. D.; Deyser, C. J.; McGowan, J. Wm.; Mitchell, J. B. A.; Wilk, S. F. J., "Merged Electron-Ion Beam Experiments I. Method and Measurements of  $(e-H_2^+)$  and  $(e-H_3^+)$  Dissociative-Recombination Cross Sections", J. Phys. B: Atom. and Molec. Phys. 10(18), 3797-820 (1977).
88. Orient, O. J., "Study of Plasma-Produced Hydrogen ( $H_3^+$ ) Ions During the Decay Period in Helium-Hydrogen Mixtures", J. Phys. B, 9(15), 2731-6 (1976).
89. Alvarez, Ignacio; Cisneros, Carmen; Barnett, C. F.; Ray, J. A., "Negative-Ion Formation From Dissociative Collisions of  $H_2^+$ ,  $H_3^+$ , and  $HD_2^+$  in Molecular Hydrogen, Helium, and Xenon", Phys. Rev. A, 14(2), 602-7 (1976).
90. Vestal, M. L.; Blakley, C. R.; Ryan, P. W.; Futrell, J. H., "Crossed-Beam Study of the Reaction  $H_3^+(D_2, H_2)D_2H^+$ ", J. Chem. Phys. 64(5), 2094-111 (1976).
91. Carney, G. D.; Porter, R. N., " $H_3^+$ : Ab Initio Calculation of the Vibration Spectrum", J. Chem. Phys. 65(9), 3547-65 (1976).
92. Johnsen, Rainer; Huang, Chou-Mou; Biondi, Manfred A., "Three-Body Association Reactions of  $H^+$  and  $H_3^+$  Ions in Hydrogen from 135 to 300 K", J. Chem. Phys. 65(4), 1539-1541 (1976).
93. Lees, A. B.; Rol, P. K., "Merging Beams Study of the  $H_3^+$  System", C. R. - Symp. Int. Jets Mol. 5th, Paper No. D<sub>3</sub>, 8 Pp.. Com. Int. Jets Mol., C/o Dr. F. Marcel Devienne: Peymeinade, Fr. (1975).
94. Anderson, James B., "Random-Walk Simulation of the Schroedinger Equation. Hydrogen Ion ( $H_3^+$ )", J. Chem. Phys. 63(4), 1499-503 (1975).
95. Hiraoka, K.; Kebarle, P., "Temperature Dependence of Third Order Ion Molecule Reactions. Reaction  $H_3^+ + 2H_2 = H_5^+ + H_2$ ", J. Chem. Phys. 63(2), 746-9 (1975).
96. Hiraoka, K.; Kebarle, P., "Determination of the Stabilities of  $H_5^+$ ,  $H_7^+$ ,  $H_9^+$ , and  $H_{11}^+$  From Measurement of the Gas Phase Ion Equilibria  $H_n^+ + H_2 = H_{n+2}^+$  ( $n = 3, 5, 7, 9$ )", J. Chem. Phys. 62(6), 2267-70 (1975).
97. Aberth, W.; Schnitzer, R.; Anbar, M., "Observations of the Diatomic and Triatomic Hydrogen Negative Ions", Phys. Rev. Lett. 34(26), 1600-03

(1975).

98. Elford, M. T.; Milloy, H. B., "Mobility of  $H_3^+$  and  $H_5^+$  Ions in Hydrogen and the Equilibrium Constant for the Reaction  $H_3^+ + 2H_2 = H_5^+ + H_2$  at Gas Temperatures of 195, 273, and 293 K", Aust. J. Phys. 27(6), 795-811 (1974).
99. Peart, B.; Dolder, K. T., "Measurements of the Dissociative Recombination of Hydrogen ( $H_3^+$ ) Ions", J. Phys. B, 7(14), 1948-52 (1974).
100. Lees, A. B.; Rol, P. K., "Merging Beams Study of the  $H_2^+(H_2, H)H_3^+$ ,  $H_2^+(D_2, H)HD_2^+$ , and  $D_2^+(H_2, H)HD_2^+$  Reaction Mechanisms", J. Chem. Phys. 61(11), 4444-9 (1974).
101. Schaad, L. J.; Hicks, W. V., "Gaussian Basis Configuration Interaction Calculations on Twenty Electronic States of  $H_3^+$ . A Bound  $^3\Sigma_u^+$  Excited Level", J. Chem. Phys. 61(5), 1934-42 (1974).
102. Bauschlicher, Charles W. Jr.; O'Neil, Stephen V.; Preston, Richard K.; Schaefer, Henry F. III, "Avoided Intersection of Potential Energy Surfaces: The ( $H^+ + H_2$ ,  $H + H_2^+$ ) System", J. Chem. Phys., 59(3), 1286-92 (1973).
103. Leu, Ming-Taun, "Recombination of Electrons with Positive Ions of the Type  $H_3^+$ ,  $H_5^+$ ,  $H_3O^+ \cdot (H_2O)_n$ , and  $HCO^+$  at Thermal Energies", 79 Pp. Avail. Univ. Microfilms, Ann Arbor, Mich., Order No. 73-13,190 From: Diss. Abstr. Int. B 1973, 33(12)(Pt. 1), 5998 (1972).
104. Patch, R. W., "Observability of the  $H_3^+$  Fundamental Spectrum", J. Chem. Phys. 57(6), 2594-5 (1972).
105. Huang, Jan-Tsyu J., "Analytical Self-Consistent-Field Energy Expression for Ground-State  $H_3^+$  Ion", J. Chem. Phys. 56(6), 3176-7 (1972).
106. Duben, Anthony J., "First Order Density Matrices From Configuration Interaction Wavefunctions for  $H_3^+$ ", 138 Pp. Avail. Univ. Microfilms, Ann Arbor, Mich., Order No. 71-16,594 From: Diss. Abstr. Int. B 1971, 32(1), 152 (1970).
107. Haung, Jan-Tsyu J., "Simple Diatomics-in-Molecules Energy Expression for  $M_3^+$  and Its Application to  $H_3^+$ ", J. Chem. Phys. 55(10), 5136-7 (1971).
108. Kawaoka, Kenji; Borkman, Raymond F., "Electric, Magnetic, and Spectral Properties of  $H_3^+$  Ground State Calculated From Single-Center Wave Functions", J. Chem. Phys. 55(9), 4637-41 (1971).
109. Middleton, C. R.; Payne, M. F.; Riviere, A. C., "Dissociation of  $H_3^+$  Ions at 410, 510, and 550 keV in Molecular Hydrogen Gas", J. Phys. B, 4(10), L88-L91 (1971).

110. Duben, Anthony J.; Lowe, John P., "Correlation Studies on  $H_3^+$ . II. Electron Densities and Expectation Values". J. Chem. Phys. 55(9), 4276-82 (1971).
111. Duben, Anthony J.; Lowe, John P., "Correlation Studies on  $H_3^+$ . I. Wave Functions", J. Chem. Phys. 55(9), 4270-5 (1971).
112. Huntress, W. T., Jr., "Ion Cyclotron Resonance Power Absorption. Collision Frequencies for  $CO_2^+$ ,  $N_2^+$ , and  $H_3^+$  Ions in Their Parent Gases", J. Chem. Phys. 55(5), 2146-55 (1971).
113. Preston, Richard K.; Tully, John C., "Effects of Surface Crossing in Chemical Reactions: The  $H_3^+$  System", J. Chem. Phys. 54(10), 4297-304 (1971).
114. Borkman, R. F., "Electric Quadrupole Moments for  $H_2^+$ ,  $H_2$ , and  $H_3^+$  from a Point-Charge Model", Chem. Phys. Lett. 9(6), 624-6 (1971).
115. Kawaoka, Kenji; Borkman, Raymond F., "Single-Center Calculations on the Electronically Excited States of Equilateral  $H_3^+$  Ion", J. Chem. Phys. 54(10), 4234-8 (1971).
116. Stecher, Theodore P.; Williams, David Arnold, "Interstellar  $H_3^+$ ", Astrophys. Lett. 7(1), 59-60 (1970).
117. Sumin, L. V.; Gur'ev, M. V., "Mechanism of Ion-Molecular Reactions of the Formation of  $H_3^+$  in Hydrogen and  $CD_5^+$  in Deuteriomethane", Dokl. Akad. Nauk SSSR, 193(4), 858-61 [Phys Chem] (1970).
118. Borkman, Raymond F., "Single-Center Configuration-Interaction Calculations on the Ground State of  $H_3^+$ ", J. Chem. Phys. 53(8), 3153-60 (1970).
119. Somorjai, R. L.; Yue, C. P., "Integral Transform Gaussian Wave Functions for  $H_3^{2+}$  and  $H_3^+$ ", J. Chem. Phys. 53(5), 1657-61 (1970).
120. Mueller, Hans, "Theoretical Study of Hydrogen Polymers. I. Calculation of the Proton Affinities of H,  $H_2$ , and  $H_3^+$ ", Z. Chem. 10(12), 478-9 (1970).
121. Csizmadia, Imre G.; Kari, R. E.; Polanyi, John C.; Roach, A. C.; Robb, M. A., "Ab-initio SCF-MO-CI Calculations for  $H^-$ ,  $H_2$ , and  $H_3^+$  Using Gaussian Basis Sets", J. Chem. Phys. 52(12), 6205-11 (1970).
122. Burt, J. A.; Dunn, Jerry L.; McEwan, M. J.; Sutton, M. M.; Roche, A. E.; Schiff, Harold I., "Ion-Molecule Reactions of  $H_3^+$  and the Proton Affinity of  $H_2$ ", J. Chem. Phys. 52(12), 6062-75 (1970).
123. Leventhal, Jacob J.; Friedman, Lewis, "Energy Transfer in the De-Excitation of  $H_3^+$ ", J. Chem. Phys. 52(12), 6062-75 (1970).

tation of  $H_3^+$  by  $H_2$ ", J. Chem. Phys. 50(7), 2928-31 (1969).

124. Preuss, H.; Janoschek, R., "Wave-Mechanical Calculations on Molecules Taking all Electrons into Account", J. Mol. Structure 3, 423-428 (1969).
125. Conroy, Harold, "Molecular Schrödinger Equation. X. Potential Surfaces for Ground and Excited States of Isosceles  $H_3^{++}$  and  $H_3^+$ ", J. Chem. Phys. 51(9), 3979-3993 (1969).
126. Poshusta, R. D.; Haugen, J. A., "Ab Initio Predictions for Very Small Ions", J. Chem. Phys. 51(8), 3343-3351 (1969).
127. Joshi, Bhairav D.; Anand, S. C., "Overlap Matrix Elements and Related Integrals Involving  $R_{12}^k$  -Type Correlation Function for  $H_3^+$  in One-Center Expansion Approximation", Indian J. Pure Appl. Phys. 6(12), 656-64 (1968).
128. Macias, A., "Configuration-Interaction Study of the  $H_3$ -System. II. Expanded Basis", J. Chem. Phys. 49(5), 2198-209 (1968).
129. Albritton, D. L.; Miller, T. M.; Martin, D. W.; McDaniel, E. W., "Mobilities of Mass-Identified  $H_3^+$  and  $H^+$  Ions in Hydrogen", Phys. Rev. 171(1), 94-102 (1968).
130. Macias, A., "Configuration-interaction Study of the  $H_3^-$  System. I. 1s Orbitals", J. Chem. Phys. 48(8), 3464-8 (1968).
131. Wu, Ay-Ju A.; Ellison, Frank O., "Method of Diatomics-in-Molecules. VII. Excited Singlet States of  $H_3^+$ ", J. Chem. Phys. 48(4), 1491-6 (1968).
132. Christoffersen, Ralph E.; Shull, Harrison, "Nature of the Two-Electron Chemical Bond. VII. Multicenter Bonds and  $H_3^+$ ", J. Chem. Phys. 48(4), 1790-7 (1968).
133. Hughes, Raymond Hargett; Kay, David B.; Stigers, C. A.; Stokes, E. D., "Production of Hydrogen Atoms in the 3s State by the Dissociation of Fast  $H_2^+$  and  $H_3^+$  Projectiles on Impact with Hydrogen, Helium, Argon, and Neon", Phys. Rev. 167(1), 26-9 (1968).
134. Neynaber, Roy H.; Trujillo, S. M., "Study of  $H_2^+ + H_2 \rightarrow H_3^+ + H$  Using Merging Beams", Phys. Rev. 167(1), 63-6 (1968).
135. Hopkinson, A. C.; Holbrook, N. K.; Yates, K.; Csizmadia, I. G., "Theoretical Study on the Proton Affinity of Small Molecules Using Gaussian Basis Sets in the LCAO-MO-SCF Framework", J. Chem. Phys. 49(8), 3596-3601 (1968).



136. Wu, Ay-Ju; Ellison, Frank O., "Method of Diatomics-in-Molecules. VIII. Excited Triplet States of  $H_3^+$ ", J. Chem. Phys. 48(11), 5032-5037 (1968).
137. Christoffersen, Ralph E.; Shull, Harrison, "Nature of the Two-Electron Chemical Bond. VII. Multicenter Bonds and  $H_3^+$ ", J. Chem. Phys. 48(4), 1790-1797 (1968).
138. Albritton, D. L.; Miller, T. M.; Moseley, J. T.; Martin, D. W.; McDaniel, E. W., "Mobilities of Mass-Identified  $H_1^+$  and  $H_3^+$  Ions in Hydrogen", Phenomena Ioniz. Gases, Int. Conf., Contrib. Pap., 8th, Vienna, 12. Springer-Verlag: Vienna, Austria. (1967).
139. Kutzelnigg, Werner; Ahlrichs, Reinhart; Labib-Iskander, I.; Bingel, Werner A., "Hartree-Fock and the Correlation Energies of the  $H_3^+$  Ion and Their Dependence on the Nuclear Configuration", Chem. Phys. Lett. 1(10), 447-50 (1967).
140. Schwartz, Maurice Edward; Schaad, Lawrence J., "Ab Initio Studies of Small Molecules Using 1s Gaussian Basis Functions. II.  $H_3^+$ ", J. Chem. Phys. 47(12), 5325-34 (1967).
141. Ritchie, Calvin D.; King, Harry Frederick, "Theoretical Studies of Proton-Transfer Reactions. I. Reactions of Hydride Ion with Hydrogen Fluoride and Hydrogen Molecules", J. Am. Chem. Soc., 90(4), 825-3 (1968).
142. Conroy, Harold; Bruner, Buddy L., "Molecular Schroedinger Equation. VI. Results for  $H_3$  and Other Simple Systems", J. Chem. Phys. 47(3), 921-9 (1967).
143. Solov'ev, E. S.; Il'in, R. N.; Oparin, V. A.; Fedorenko, N. V., "Production of Highly Excited Hydrogen Molecules and Atoms by Fast  $H_2^+$  and  $H_3^+$  Ions Passing Through Hydrogen, Neon, and Magnesium and Sodium Vapor", Zh. Eksp. Teor. Fiz. 53(6), 1933-41 (1967).
144. Devienne, F. Marcel; Roustau, Jean C., "Existence and Some Properties of Triatomic Molecular Jets of Hydrogen", C. R. Acad. Sci., Ser. A B (See CHASAP and CHDBAN), 263B(25), 1389-92 (1966).
145. Pfeiffer, Gary V.; Huff, Norman T.; Greenawalt, E. M.; Ellison, Frank O., "Method of Diatomics in Molecules. IV. Ground and Excited States of  $H_3^+$ ,  $H_4^+$ ,  $H_5^+$  and  $H_6^+$ ", J. Chem. Phys. 46(2), 821-2 (1967).
146. Considine, James P.; Hayes, Edward F., "Single-Center Wave Functions for  $H_3^+$  and  $H_3$ ", J. Chem. Phys. 46(3), 1119-24 (1967).
147. Wu, Ay-Ju; Ellison, Frank O., "Method of Diatomics-in-Molecules. VII. Excited Singlet States of  $H_3^+$ ", J. Chem. Phys. 48(4), 1491-1497 (1967).

148. Kutzelnigg, W.; Ahlrichs, R.; Labib-Iskander, I.; Bingel, W. A., "The Hartree-Fock and the Correlation Energies of the  $H_3^+$  Ion and Their Dependence on the Nuclear Configuration", Chem. Phys. Lett. 1(10), 447-450 (1967).
149. Schwartz, Maurice E.; Schaad, L. J., "Ab Initio Studies of Small Molecules Using 1s Gaussian Basis Functions. II.  $H_3^+$ ", J. Chem. Phys. 47(12), 5325-5334 (1967).
150. Considine, J. P.; Hayes, E. F., "Single-Center Wavefunctions for  $H_3^+$  and  $H_3$ ", J. Chem. Phys. 46(3), 1119-1124 (1967).
151. Buchheit, K.; Henkes, W., "Untersuchung der Massenverteilung von Wasserstoff-Agglomerat-Ionen in Einem Massenspektrometer mit Energiezerlegung", Z. Angew. Phys. 24(4), 191-6 (1967).
152. Schwartz, Maurice Edward, "Hellmann-Feynman Theorem and the Correlation Energy of  $H_3^+$ ", J. Chem. Phys. 45(12), 4754-5 (1966).
153. Williams, James Francis; Dunbar, D. N. F., "Charge Exchange and Dissociation Cross Sections for  $H_1^+$ ,  $H_2^+$ , and  $H_3^+$  Ions of 2- to 50-keV Energy Incident Upon Hydrogen and the Inert Gases", Phys. Rev. 149(1), 62-9 (1966).
154. Joshi, Bhairav D., "Study of the  $H_3^+$  Molecule Using Self-Consistent-Field One-Center Expansion Approximation", J. Chem. Phys. 44(9), 3627-3631 (1966).
155. Pearson, A. G.; Poshusta, R. D.; Browne, J. C., "Some Potential-Energy Surfaces on  $H_3^+$  Computed with Generalized Gaussian Orbitals", J. Chem. Phys. 44(5), 1815-1818 (1966).
156. Lester, William A. Jr.; Krauss, Morris, "Some Aspects of the Coulomb Hole of the Ground State of  $H_3^+$ ", J. Chem. Phys., 44(1), 207-212 (1966).

Section III:  $H_N^+$  (and  $H_N^-$ ,  $N \geq 4$ )

1. Hobza, P.; Schneider, B.; Sauer, J.; Carsky, P.; Zahradnik, R., "MP4 Interaction Energies and Basis Set Superposition Errors for the Molecular Hydrogen ( $(H_2)_2$ ) Dimer", Chem. Phys. Lett. 134(5), 418-22 (1987).
2. Melnikopoulos, A.; Nicolaides, C. A., "Towards Understanding the Stability of the  $H_4^+$  ( $C_{3v}$ ) Cluster", Z. Phys. D: At., Mol. Clusters, 5(2), 175-80 (1987).
3. Theodorakopoulos, G.; Petsalakis, I. D.; Nicolaides, C. A., "Potential Energy Hypersurfaces of the Hydrogen Tetraatomic Molecule in the Ground and the First Two Singlet Excited Electronic States", THEOCHEM, 34(1-2), 23-31 (1987).
4. Yamaguchi, Yukio; Gaw, Jeffrey F.; Remington, Richard B.; Schaefer, Henry F. III; "The  $H_5^+$  Potential Energy Hypersurface: Characterization of Ten Distinct Energetically Low-Lying Stationary Points", J. Chem. Phys. 86(9), 5072-5081 (1987).
5. Hiraoka, Kenzo, "A Determination of the Stabilities of  $H_3^+(H_2)_n$  with  $n = 1-9$  from Measurements of the Gas-Phase Ion Equilibria  $H_3^+(H_2)_{n-1} + H_2 = H_3^+(H_2)_n$ ", J. Chem. Phys. 87(7), 4048-55 (1987).
6. Kirchner, N. J.; Bowers, M. T., "An Experimental Study of the Formation and Reactivity of Ionic Hydrogen Clusters: The First Observation and Characterization of the Even Clusters  $H_4^+$ ,  $H_6^+$ ,  $H_8^+$ , and  $H_{10}^+$ ", J. Chem. Phys. 86(3), 1301-1310 (1987).
7. Kirchner, N. J.; Bowers, M. T., "Fragmentation Dynamics of Metastable Hydrogen Ion Clusters  $H_5^+$ ,  $H_7^+$  and  $H_9^+$ : Experiment and Theory", J. Phys. Chem. 91, 2573-2582 (1987).
8. Forward, R., "Prospects for Antiproton Production and Propulsion", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 9-26. Edited by John T. Bahns. University of Dayton Research Institute (1987).
9. Saxon, R., "Overview of Hydrogen Clusters", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 27-37. Edited by John T. Bahns. University of Dayton Research Institute (1987).
10. Stwalley, W. C., "Large Hydrogen Cluster Ions", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 39-45. Edited by John T. Bahns. University of Dayton Research Institute (1987).
11. Bahns, J. T., "Introduction to the Workshop on Cooling, Condensation and

Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 1-7. Edited by John T. Bahns. University of Dayton Research Institute (1987).

12. Bowers, M. T., "Formation and Reactivity of Small Hydrogen Cluster Ions", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 61-72. Edited by John T. Bahns. University of Dayton Research Institute (1987).
13. Yamaguchi, Yukio; Gaw, Jeffrey F.; Remington, Richard J.; Schaefer, Henry F. III, "The  $H_5^+$  Potential Energy Hypersurface. Characterization of Ten Distinct Energetically Low-Lying Stationary Points", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 73-79. Edited by John T. Bahns. University of Dayton Research Institute (1987).
14. Montgomery, J. A. Jr.; Michels, H. H., "Electronic Structure and Stability of Small Cation and Anion Hydrogen Clusters", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 81-94. Edited by John T. Bahns. University of Dayton Research Institute (1987).
15. Crofton, M., "Infrared Spectroscopy of  $H_3^+$ ", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 119-124. Edited by John T. Bahns. University of Dayton Research Institute (1987).
16. Yeh, L. I.; Okumura, M.; Myers, J. D.; Lee, Y. T., "Vibrational Spectroscopy of the Hydrogen and Hydrated Hydronium Cluster Ions", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 125-141. Edited by John T. Bahns. University of Dayton Research Institute (1987).
17. Mitchell, J. B. A., "The Role of Electron-Ion Recombination in Bulk Antimatter Production", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 143-156. Edited by John T. Bahns. University of Dayton Research Institute (1987).
18. Bahns, J. T., "Key Problems and Hydrogen Atom Formation", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, 219-230. Edited by John T. Bahns. University of Dayton Research Institute (1987).
19. Bahns, J. T.; Sando, K. M.; Tardy, D. C.; Stwalley, W. C., "Proceedings of the Hydrogen Cluster Ion Study Group", Proc. Cooling, Condensation, and Storage of Hydrogen Cluster Ions Workshop, 1/8-9/87, A1-64. Edited by John T. Bahns. University of Dayton Research Institute (1987).
20. Cardelino, B. H.; Eberhardt, W. H.; Borkman, R. F., "Ab Initio SCF Calculation on Lithium-Hydrogen ( $Li_nH_m$ ) Molecules and Cations with Four or Less Atoms", J. Chem. Phys. 84(6), 3230-42 (1986).

21. Roszak, S.; Sokalski, W. A.; Hariharan, P. C.; Kaufman, Joyce J., "Procedure Supplementing SCF Interaction Energies by Dispersion Term Evaluated in Dimer Basis Set within Variation-Perturbation Approach", *Theor. Chim. Acta*, 70(2), 81-8 (1986).
22. Yeh, L. I.; Okumura, M.; Lee, Y. T., "The Vibrational Predissociation Spectroscopy of Hydrogen Cluster Ions", 813-818 in "Electronic and Atomic Collisions", Edited by D. C. Lorents, W. E. Meyerhof, J. R. Peterson, Elsevier Science Publishers B. V., NY (1986).
23. Okumura, M.; Yeh, L. I.; Lee, Y. T., "The Vibrational Predissociation Spectroscopy of Hydrogen Cluster Ions", *J. Chem. Phys.*, 83(7), 3705-6 (1985).
24. Nicolaidis, C. A.; Theodorakopoulos, G.; Petsalakis, I. D., "Theory of Chemical Reactions of Vibronically Excited Diatomic Hydrogen ( $B^1\Sigma_u^+$ ). I. Prediction of a Strongly Bound Excited State of Tetraatomic Hydrogen", *J. Chem. Phys.* 80(4), 1705-6 (1984).
25. Kirchner, Nicholas J.; Gilbert, James R.; Bowers, Michael T., "The First Experimental Observation of Stable Hydrogen ( $H_4^+$ ) Ions", *Chem. Phys. Lett.* 106(1-2), 7-12 (1984).
26. MacDonald, Jeffrey A.; Biondi, Manfred A.; Johnsen, Rainer, "Recombination of Electrons with Hydrogen ( $H_3^+$  and  $H_5^+$ ) Ions", *Planet. Space Sci.* 32(5), 651-4 (1984).
27. Beuhler, R. J.; Friedman, L., "Cluster Ion Formation in Free Jet Expansion Process at Low Temperatures", *Ber. Bunsenges. Phys. Chem.* 88, 265 (1984).
28. Moser, H. O., "Time of Flight Spectroscopy of Sub-meV Cluster Ions in the Mass Range  $1-10^6$  Atoms per Charge", *Rev. Sci. Instrum.* 55(12), 1914-1923 (1984).
29. Kutina, R. E.; Edwards, A. K.; Pandolf, R. S.; Berkowitz, J., "UV Laser Photodissociation of Molecular Ions", *J. Chem. Phys.* 80(9), 4112-9 (1984).
30. Jungen, Martin; Staemmler, Volker, "Rydberg States of the Hydrogen Tetramer ( $H_4$ )", *Chem. Phys. Lett.* 103(3), 191-5 (1983).
31. Pulay, Peter, "Variational Formulation and Gradient Evaluation for Coupled Electron Pair Approximations: A Model Study", *Int. J. Quantum Chem., Quantum Chem. Symp.* 17, 257-63 (1983).
32. Takahashi, Mitsuo; Fukutome, Hideo, "Projected BCS Tamm-Dancoff Method for Molecular Electronic Structures", *Int. J. Quantum Chem.* 24(6), 603-21 (1983).

33. Brown, Richard E.; Colpa, Johannes Pieter, "SCF and CI Studies of Hund's Rules for the Effects of Electronic Correlation and Delocalization. I", *Int. J. Quantum Chem.* 24(6), 593-602 (1983).
34. Wilson, S.; Jankowski, K.; Paldus, J., "Applicability of Nondegenerate Many-Body Perturbation Theory to Quasi-Degenerate Electronic States: A Model Study", *Int. J. Quantum Chem.* 23(5), 1781-802 (1983).
35. Yamaguchi, Yukio; Gaw, Jeffrey F.; Schaefer, Henry F., III, "Molecular Clustering about a Positive Ion. Structures, Energetics, and Vibrational Frequencies of the Protonated Hydrogen Clusters  $H_3^+$ ,  $H_5^+$ ,  $H_7^+$ , and  $H_9^+$ ", *J. Chem. Phys.* 78(6, Pt. 2), 4074-85 (1983).
36. Beuhler, R. J.; Ehenson, S.; Friedman, L., "Hydrogen Cluster Ion Equilibria", *J. Chem. Phys.* 79(12), 5982-90 (1983).
37. Elford, M. T., "The Heat of Dissociation of  $H_5^+$  Derived from Measurements of Ion Mobilities", *J. Chem. Phys.* 79(12) 5951-5959 (1983).
38. Hirao, K.; Yamabe, S., "Theoretical Study on the Structure and Stability of Hydrogen Ion Clusters  $H_n^+$  and  $H_n^-$  ( $n = 3, 5, 7, 9, 11, 13$ )", *Chem. Phys.* 80, 237-243 (1983).
39. Huber, H.; Szekeley, D., "Near-Hartree-Fock Energies and Geometries of the Hydrogen Clusters  $H_n^+$  ( $n$  (odd)  $\leq 13$ ) Obtained with Floating Basis Sets", *Theor. Chim. Acta* 62, 499-506 (1983).
40. Raynor, S.; Herschbach, D. R., "Electronic Structure of  $H_n^+$  and  $HeH_n^+$  Clusters", *J. Phys. Chem.* 87, 289-293 (1983).
41. Wright, L. R.; Borkman, R. F., "Ab Initio Studies on the Stabilities of Even- and Odd-Membered Hydrogen ( $H_n^+$ ) Clusters", *J. Chem. Phys.* 77(4), 1938-41 (1982).
42. Borisov, Yu. A., "The Functional of the Electron Density for Atoms and Molecules", *Chem. Phys. Lett.* 93(2), 197-200 (1982).
43. Beuhler, R. J.; Friedman, L., "Hydrogen Cluster Ions", *Phys. Rev. Lett.* 48(16), 1097-99 (1982).
44. Chanut, Y.; Martin, J.; Salin, R.; Moser, H.O., "Production and Beam Analysis of Energetic Small Hydrogen Cluster Ions for Study of Their Interactions with Targets and Their Structure", *Surface Science* 106(1-3), 563-8 (1981).
45. Yano, K.; Bee, S. H., "Mass Analyses of Cluster Ion Beams by Wein Filter", *Jap. J. of Appl. Phys.* 19(6), 1019-25 (1980).
46. Huber, H., "Geometry Optimization in Ab Initio SCF Calculations. The Hydrogen Clusters  $H_n^+$  ( $n = 7, 9, 11, 13$ )", *Chem. Phys. Lett.* 70(2), 353-7

(1980).

47. Anderson, S. L.; Hirooka, T.; Tiedelmann, P. W.; Mahan, B. H.; Lee, Y. T., "Photoionization of  $(H_2)_2$  and Clusters of  $O_2$  Molecules", J. Chem. Phys. 73(10), 4779-4783 (1980).
48. Bunker, P. R., "Symmetry in Molecular Hydrogen Dimer  $((H_2)_2)$ , Molecular Deuterium Dimer  $((D_2)_2)$ , Hydrogen Deuteride Dimer  $((HD)_2)$ , and  $H_2$ - $D_2$  Van der Waals Complexes", Can. J. Phys. 57(12), 2099-105 (1979).
49. Vojtik, J.; Polak, R., "On the Multidimensional "Avoided Surface Crossing" Problem", Chem. Phys. 42(1-2), 177-82 (1979).
50. Van Lumig, A.; Reuss, J.; Ding, A.; Weise, J.; Rindtisch, A., "Double Differential Fragmentation Cross Section Measurements of  $H_{2n+1}^+$  Ions,  $n \leq 7$ ", Mol. Phys. 38(2), 337-351 (1979).
51. Huber, H., "Geometry Optimization in Ab Initio SCF Calculations. Floating Orbital Geometry Optimization Applying the Hellmann-Feynman Force", Chem. Phys. Lett. 62(1), 95-99 (1979).
52. Sapse, A. M.; Rayez-Meaume, M. T.; Rayez, J. C.; Massa, L. J., "Ion-Induced Dipole  $H_n^-$  Clusters", Nature 278, 332-33 (1979).
53. Van Lumig, A.; Reuss, J., "Collisions of Hydrogen Cluster Ions with a Gas Target, at 200-850 eV Energy", Int. J. Mass Spectrom. Ion Phys. 27(2), 197-208 (1978).
54. Polak, R.; Vojtik, J.; Schneider, F., "Analysis of DIM Energy Hypersurfaces for Some Hydrogenic and Lithium Clusters", Chem. Phys. Lett. 53(1), 117-20 (1978).
55. Stine, J. R.; Muckerman, J. T., "Charge Exchange and Chemical Reaction in the  $H_2^+ + H_2$  System. I. Characterization of the Potential Energy Surfaces and Nonadiabatic Regions", J. Chem. Phys. 68(1), 185-94 (1978).
56. Polak, R., "Diatomics-in-Molecules Study on the Stability of the  $H_4^+$  Ion", Chem. Phys. 16(3), 353-9 (1976).
57. Krenos, J. R.; Lehmann, K. K.; Tully, J. C.; Hierl, P. M.; Smith, G. P., "Crossed-Beam Study of the Reactions of Molecular Hydrogen(+) with Molecular Deuterium and Molecular Deuterium(+) with Molecular Hydrogen", Chem. Phys. 16(1), 109-16 (1976).
58. Arifov, U. A.; Lugovskoi, V. B.; Makarenko, V. A., "Emission of Negative Ions During Irradiation of Metals by Microsecond Pulses of Ruby Laser Radiation", Izv. Akad. Nauk SSSR, Ser. Fiz. 40(8), 1702-6 (1976).
59. Cobb, M.; Moran, T. F.; Borkman, R. F.; Childs, R., "Ab Initio Potential Energy Curves for  $H_2$ - $H_2^+$  Interactions", Chem. Phys. Lett. 57(3), 326-30

(1978).

60. Yamabe, S.; Hirao, K.; Kitaura, K., "Theoretical Study on the Stability and the Structure of  $H_n^+$  ( $n = 3, 5, 7, 9, 11$ )", Chem. Phys. Lett. 56(3), 546-8 (1978).
61. Johnsen, Rainer; Huang, Chou-Mou; Biondi, Manfred A., "Three-Body Association Reactions of  $H^+$  and  $H_3^+$  Ions in Hydrogen from 13' to 300 K", J. Chem. Phys. 65(4), 1539-1541 (1976).
62. Carsky, Petr; Zahradnik, Rudolf; Hobza, Pavel, "Semiempirical Estimates of the Correlation Energy in Small Clusters of Hydrogen Atoms", Theor. Chim. Acta, 40(4), 287-95 (1975).
63. Hiraoka, K.; Kebarle, P., "Temperature Dependence of Third Order Ion Molecule Reactions. Reaction  $H_3^+ + 2H_2 = H_5^+ + H_2$ ", J. Chem. Phys. 63(2), 746-9 (1975).
64. Hiraoka, K.; Kebarle, P., "Determination of the Stabilities of  $H_5^+$ ,  $H_7^+$ ,  $H_9^+$ , and  $H_{11}^+$  From Measurement of the Gas Phase Ion Equilibria  $H_n^+ + H_2 = H_{n+2}^+$  ( $n = 3, 5, 7, 9$ )", J. Chem. Phys. 62(6), 2267-70 (1975).
65. Aberth, W.; Schnitzer, R.; Anbar, M., "Observations of the Diatomic and Triatomic Hydrogen Negative Ions", Phys. Rev. Lett. 34(26), 1600-03 (1975).
66. Ahlrichs, R., "Theoretical Study of the  $H_5^+$  System", Theoret. Chim. Acta 39, 149-160 (1975).
67. Elford, M. T.; Milloy, H. B., "Mobility of  $H_3^+$  and  $H_5^+$  Ions in Hydrogen and the Equilibrium Constant for the Reaction  $H_3^+ + 2H_2 = H_5^+ + H_2$  at Gas Temperatures of 195, 273, and 293 K", Aust. J. Phys. 27(6), 795-811 (1974).
68. Harrison, S. W.; Massa, L. J.; Solomon, P., "Binding Energy and Geometry of the Hydrogen Clusters  $H_n^+$ ", Nature 245, 31-2 (1973).
69. Salmon, W. I.; Poshusta, R. D., "Polarized-Orbital Valence-Bond Calculations on the Ground State Properties of  $H_5^+$ ", J. Chem. Phys. 59(9), 4867-70 (1973).
70. Van Deursen, A.; Reuss, J., "Measurements of Intensity and Velocity Distribution of Clusters from a  $H_2$  Supersonic Nozzle Beam", Int. J. of Mass Spectrom. and Ion Phys. 11, 483-489 (1973).
71. Leu, Ming-Taun, "Recombination of Electrons with Positive Ions of the Type  $H_3^+$ ,  $H_5^+$ ,  $H_3O^+ \cdot (H_2O)_n$ , and  $HCO^+$  at Thermal Energies", 79 Pp. Avail. Univ. Microfilms, Ann Arbor, Mich., Order No. 73-13,190 From: Diss. Abstr. Int. B 1973, 33(12)(Pt. 1), 5998 (1972).



72. Bennett, S. L.; Field, F. H.; "Reversible Reactions of Gaseous Ions. VII. The Hydrogen System", J. Am. Chem. Soc. 94(25), 8669-72 (1972).
73. Andreev, St., "Configurations of the Polyatomic Hydrogen Molecules  $H_4^+$  and  $H_5^+$ ", God. Vissh. Khimikotekhnol. Inst., Sofia, Volume Date 1967, 14(1), 85-90 (1971).
74. Arifov, U. A.; Pozharov, S. L.; Chernov, I. G.; Mukhamediev, Z. A., "Ionic-Molecular Reactions in Hydrogen at High Pressures", High Energy Chemistry 5, 69 (1971).
75. Clampitt, R.; Gowland, L., "Clustering of Cold Hydrogen Gas on Protons", Nature 223, 815-816 (1969).
76. Schwartz, M. E.; Schaad, L. J., "Ab Initio Studies of Small Molecules Using 1s Gaussian Basis Functions. III. LCGTO SCF MO Wavefunctions of the Three- and Four-Electron Systems  $He_2^+$ ,  $He_2$ , and Linear  $H_3$ ,  $H_4^+$ ,  $H_4$ ", J. Chem. Phys. 48(10), 4709-15 (1968).
77. Pfeiffer, Gary V.; Huff, Norman T.; Greenawalt, E. M.; Ellison, Frank O., "Method of Diatomics in Molecules. IV. Ground and Excited States of  $H_3^+$ ,  $H_4^+$ ,  $H_5^+$  and  $H_6^+$ ", J. Chem. Phys. 46(2), 821-2 (1967).
78. Buchheit, K.; Henkes, W., "Untersuchung der Massenverteilung von Wasserstoff-Agglomerat-Ionen in Einem Massenspektrometer mit Energiezerlegung", Z Angew. Phys. 24(4), 191-6 (1967).